



US012076198B2

(12) **United States Patent**
Colby

(10) **Patent No.:** **US 12,076,198 B2**
(45) **Date of Patent:** **Sep. 3, 2024**

(54) **THERAPEUTIC TOOTH BUD ABLATION**
(71) Applicant: **TriAgenics, Inc.**, Redmond, OR (US)
(72) Inventor: **Leigh E. Colby**, Lino Lakes, MN (US)
(73) Assignee: **TRIAGENICS, INC.**, Redmond, OR (US)
(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 1401 days.

(51) **Int. Cl.**
A61C 1/08 (2006.01)
A61C 3/00 (2006.01)
A61C 5/42 (2017.01)
(52) **U.S. Cl.**
CPC *A61C 1/082* (2013.01); *A61C 3/00* (2013.01); *A61C 5/42* (2017.02); *Y10T 29/49826* (2015.01)

(21) Appl. No.: **16/391,277**

(22) Filed: **Apr. 22, 2019**

(65) **Prior Publication Data**
US 2019/0247144 A1 Aug. 15, 2019

Related U.S. Application Data

(63) Continuation of application No. 15/886,729, filed on Feb. 1, 2018, now Pat. No. 10,299,885, which is a continuation of application No. 15/694,794, filed on Sep. 2, 2017, now Pat. No. 10,285,778, which is a continuation of application No. 13/093,842, filed on Apr. 26, 2011, now Pat. No. 9,827,068, which is a continuation of application No. 12/863,183, filed as application No. PCT/US2010/034259 on May 10, 2010, now Pat. No. 9,402,693, application No. 16/391,277 is a continuation of application No. 15/694,794, filed on Sep. 2, 2017, now Pat. No. 10,285,778, and a continuation of application No. 15/694,791, filed on Sep. 2, 2017, now Pat. No. 10,265,140, and a continuation of application No. 13/093,844, filed on Apr. 26, 2011, now Pat. No. 10,820,963.

(60) Provisional application No. 61/177,143, filed on May 11, 2009.

(58) **Field of Classification Search**
CPC .. *A61C 1/082*; *A61C 1/00*; *A61C 5/42*; *A61C 3/00*; *A61C 5/44*; *A61C 5/46*; *Y10T 29/49826*
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

610,355 A 9/1898 Kinne et al.
752,378 A 2/1904 Dailey
1,713,971 A 5/1929 Lowry et al.
(Continued)

FOREIGN PATENT DOCUMENTS

AU 2010247874 11/2010
AU 2015213338 9/2015
(Continued)


OTHER PUBLICATIONS

X-Nav Technologies; "How It Works: X-Guide Dynamic 3D Navigation™ Workflow;" X-Nav Technologies; 2 pages; at least as early as 2020. <http://www.x-navtech.com/x-guide/how-it-works>.
(Continued)

Primary Examiner — Yogesh P Patel
(74) *Attorney, Agent, or Firm* — LAW OFFICE OF KAREN DANA OSTER, LLC

(57) **ABSTRACT**
A custom surgical stent for use in a tooth bud ablation procedure that results in tooth agenesis.

6 Claims, 29 Drawing Sheets



(56)

References Cited

U.S. PATENT DOCUMENTS

2,317,648 A	4/1943	Siqveland	7,812,815 B2	10/2010	Banerjee et al.
4,666,130 A	5/1987	Denman	7,819,591 B2	10/2010	Rohaly et al.
4,672,963 A	6/1987	Barken	7,822,466 B2	10/2010	Stoisnovici et al.
4,700,716 A	10/1987	Kasevich et al.	7,959,441 B2	6/2011	Glover et al.
4,895,517 A	1/1990	Fischer	7,976,469 B2	7/2011	Bonde et al.
4,925,523 A	5/1990	Braren et al.	7,990,548 B2	8/2011	Babayoff et al.
5,009,595 A	4/1991	Osborn	8,013,853 B1	9/2011	Douglas et al.
5,015,183 A	5/1991	Fenick	8,048,067 B2	11/2011	Davalos et al.
5,028,463 A	7/1991	Cahill	8,057,487 B2	11/2011	Chu et al.
5,133,660 A	7/1992	Fenick	8,211,099 B2	7/2012	Buysse et al.
5,261,647 A	11/1993	Venegas, Jr.	8,221,121 B2	7/2012	Berckmans et al.
5,443,324 A	8/1995	Sullivan	8,257,341 B1	9/2012	Fletcher
5,480,417 A	1/1996	Hascoet	8,310,683 B2	11/2012	Babayoff et al.
5,500,012 A	3/1996	Brucker et al.	8,363,228 B2	1/2013	Babayoff
5,556,278 A	9/1996	Meitner	8,372,061 B2	2/2013	Berna et al.
5,613,852 A	3/1997	Bavitz	8,377,057 B2	2/2013	Rick et al.
5,623,931 A	4/1997	Wung et al.	8,417,010 B1	4/2013	Colby
5,642,997 A	7/1997	Gregg, II et al.	8,480,666 B2	7/2013	Buysse et al.
5,688,118 A	11/1997	Hayka et al.	8,626,317 B2	1/2014	Higgins
5,725,376 A	3/1998	Poirier	8,670,816 B2	3/2014	Green et al.
5,768,134 A	6/1998	Swaelens et al.	8,672,932 B2	3/2014	Van Der Weide
5,792,140 A	8/1998	Tu et al.	8,807,864 B2	8/2014	Kulkarni
5,800,168 A	9/1998	Cascione et al.	8,834,409 B2	9/2014	Manley
5,816,814 A	10/1998	Venta et al.	8,892,235 B2	11/2014	Choi et al.
5,842,858 A	12/1998	Truppe	8,945,144 B2	2/2015	Cunningham
5,843,021 A	12/1998	Edwards et al.	8,979,871 B2	3/2015	Tyc et al.
5,851,112 A	12/1998	Daikuzono et al.	9,028,252 B1	5/2015	Shabat
5,876,020 A	3/1999	Giavotto	9,066,712 B2	6/2015	Fourkas et al.
5,906,609 A	5/1999	Assa et al.	9,072,532 B2	7/2015	Van Der Weide
5,916,213 A	6/1999	Haissaguerre	9,113,912 B1	8/2015	Mehta et al.
5,941,845 A	8/1999	Tu et al.	9,119,628 B1	9/2015	Mehta et al.
5,941,889 A	8/1999	Cermak	9,119,649 B2	9/2015	Van Der Weide
5,943,719 A	8/1999	Feldman et al.	9,168,091 B2	10/2015	Janssen et al.
5,967,777 A	10/1999	Klein et al.	9,192,438 B2	11/2015	Thiel
6,027,497 A	2/2000	Daniel et al.	9,271,796 B2	3/2016	Buysse et al.
6,149,134 A	11/2000	Bank	9,339,245 B2	5/2016	Colby
6,162,052 A	12/2000	Kokubu	9,402,691 B2	8/2016	Merritt et al.
6,179,803 B1	1/2001	Edwards et al.	9,402,693 B2	8/2016	Colby
6,179,830 B1	1/2001	Kokubu	9,415,194 B2	8/2016	Wolf et al.
6,203,499 B1	3/2001	Imling et al.	9,422,159 B2	8/2016	Colby
6,210,355 B1	4/2001	Edwards et al.	9,425,234 B2	8/2016	Colby
6,241,425 B1	6/2001	Kazim	9,439,608 B2	9/2016	Schutysen et al.
6,241,725 B1	6/2001	Cosman	9,566,115 B2	2/2017	Van Der Weide
6,245,062 B1	6/2001	Berube et al.	9,597,160 B1	3/2017	Gregg, II
6,270,476 B1	8/2001	Santojanni et al.	9,622,813 B2	4/2017	Krugman et al.
6,319,006 B1	11/2001	Scherer et al.	9,724,236 B2	8/2017	Abe
6,386,869 B1	5/2002	Zegarelli	9,827,068 B2	11/2017	Colby
6,607,524 B1	8/2003	LaBudde et al.	9,839,402 B2	12/2017	Colby
6,621,491 B1	9/2003	Baumrind et al.	9,844,324 B2	12/2017	Merritt et al.
6,622,731 B2	9/2003	Daniel et al.	9,855,112 B2	1/2018	Colby
6,913,463 B2	7/2005	Blacklock	9,861,440 B2	1/2018	Van Der Weide
6,936,046 B2	8/2005	Hissong et al.	9,872,729 B2	1/2018	Van Der Weide
6,971,877 B2 *	12/2005	Harterer A61C 1/084 433/75	9,877,783 B2	1/2018	Van Der Weide
7,014,461 B2	3/2006	Weinstein	9,943,374 B2	4/2018	Merritt et al.
7,044,735 B2	5/2006	Malin	9,943,379 B2	4/2018	Gregg, II
7,097,451 B2	8/2006	Tang	10,022,202 B2	7/2018	Colby
7,169,155 B2	1/2007	Chu et al.	10,265,140 B2	4/2019	Colby
7,226,288 B2	6/2007	Schoeffel	10,285,778 B2	5/2019	Colby
7,249,952 B2	7/2007	Ranta et al.	10,298,255 B2	5/2019	Colby
7,346,417 B2	3/2008	Luth et al.	10,299,885 B2	5/2019	Colby
7,445,402 B1	11/2008	Chen	10,330,401 B2	6/2019	Clarke
7,457,443 B2	11/2008	Persky	10,335,248 B2	7/2019	Colby
7,467,015 B2	12/2008	van der Weide	10,350,008 B2	7/2019	Gibbs et al.
7,492,987 B2	2/2009	Yeik et al.	10,355,248 B2	7/2019	Colby
7,520,877 B2	4/2009	Lee, Jr. et al.	10,765,490 B2	9/2020	Colby
7,553,309 B2	6/2009	Buysse et al.	10,820,963 B2	11/2020	Colby
7,565,190 B2	7/2009	Okerlund et al.	11,173,012 B2	11/2021	Colby
7,611,508 B2	11/2009	Yang	11,399,915 B2	8/2022	Colby
7,615,047 B2	11/2009	Berna et al.	11,540,881 B2	1/2023	Cao
7,725,151 B2	5/2010	Warren	11,583,337 B2	2/2023	Colby
7,736,357 B2	6/2010	Lee, Jr. et al.	11,864,961 B2	1/2024	Colby
7,736,537 B1	6/2010	Zastrow	2002/0022864 A1	2/2002	Mahvi
7,776,035 B2	8/2010	Rick et al.	2002/0058872 A1	5/2002	Steininger et al.
7,787,132 B2	8/2010	Körner et al.	2002/0128642 A1	9/2002	Berube et al.
			2002/0160337 A1	10/2002	Klein et al.
			2003/0130575 A1	7/2003	Desai
			2004/0097805 A1	5/2004	Verard et al.
			2004/0102767 A1	5/2004	Stingl et al.
			2004/0147915 A1	7/2004	Hasebe

(56)

References Cited

U.S. PATENT DOCUMENTS

2004/0157188 A1 8/2004 Luth et al.
 2004/0158194 A1 8/2004 Wolff et al.
 2004/0215186 A1 10/2004 Cornelius
 2004/0219482 A1 11/2004 Bina et al.
 2004/0230187 A1 11/2004 Lee
 2004/0249370 A1 12/2004 Berna et al.
 2005/0015081 A1 1/2005 Turovskiy
 2005/0090820 A1 4/2005 Cornelius
 2005/0170312 A1 8/2005 Pond
 2005/0177074 A1 8/2005 Becker et al.
 2005/0186533 A1 8/2005 Cohen
 2005/0245817 A1 11/2005 Clayton
 2005/0283148 A1 12/2005 Janssen et al.
 2006/0009404 A1 1/2006 Williams
 2006/0074413 A1 4/2006 Behzadian
 2006/0084958 A1 4/2006 Raif et al.
 2006/0111705 A1 5/2006 Janzen et al.
 2006/0115782 A1 6/2006 Li et al.
 2006/0125572 A1 6/2006 Dan Der Weide
 2006/0127859 A1 6/2006 Wen
 2006/0195169 A1 8/2006 Gross et al.
 2006/0240378 A1 10/2006 Weinstein et al.
 2006/0265139 A1 11/2006 Dan Der Weide
 2006/0276781 A1 12/2006 Dan Der Weide
 2006/0278825 A1 12/2006 Dan Der Weide
 2007/0016181 A1 1/2007 Dan Der Weide
 2007/0016183 A1 1/2007 Lee
 2007/0049917 A1 3/2007 Yang et al.
 2007/0049918 A1 3/2007 Dan Der Weide
 2007/0055224 A1 3/2007 Lee, Jr.
 2007/0135707 A1 6/2007 Redel et al.
 2007/0179485 A1 8/2007 Yeik et al.
 2007/0203551 A1 8/2007 Cronin et al.
 2007/0224570 A1 9/2007 West et al.
 2007/0265628 A1 11/2007 Kraus et al.
 2007/0288079 A1 12/2007 Dan Der Weide
 2007/0293756 A1 12/2007 Jung et al.
 2007/0294280 A1 12/2007 Jung et al.
 2008/0009747 A1 1/2008 Saadat
 2008/0033422 A1 2/2008 Turner et al.
 2008/0033424 A1 2/2008 Dan Der Weide
 2008/0039709 A1 2/2008 Karmarkar
 2008/0045938 A1 2/2008 Dan Der Weide
 2008/0119921 A1 5/2008 Brace
 2008/0138761 A1 6/2008 Pond
 2008/0147056 A1 6/2008 Dan Der Weide
 2008/0176187 A1 7/2008 Stumpel
 2008/0182218 A1 7/2008 Chen et al.
 2008/0199829 A1 8/2008 Paley et al.
 2008/0275436 A1 11/2008 Cronin
 2009/0011382 A1 1/2009 Bavar
 2009/0041198 A1 2/2009 Price
 2009/0062790 A1 3/2009 Malchano
 2009/0076498 A1 3/2009 Saadat et al.
 2009/0088830 A1 4/2009 Mohamed et al.
 2009/0118613 A1 5/2009 Krugman et al.
 2009/0136902 A1 5/2009 Zundorf et al.
 2009/0253095 A1 10/2009 Salcedo et al.
 2009/0258330 A1 10/2009 Huber et al.
 2009/0263764 A1 10/2009 Berckmans et al.
 2009/0306652 A1 12/2009 Buysse
 2010/0021866 A1 1/2010 Tsuji et al.
 2010/0030061 A1 2/2010 Canfield
 2010/0035201 A1 2/2010 Beck et al.
 2010/0082032 A1 4/2010 Berna et al.
 2010/0100094 A1 4/2010 Truckai
 2010/0114184 A1 5/2010 Degtyar et al.
 2010/0129771 A1 5/2010 Tsuji et al.
 2010/0203479 A1 8/2010 Bulloch et al.
 2010/0268219 A1 10/2010 Ornsby et al.
 2010/0298705 A1 11/2010 Pelissier
 2010/0311006 A1 12/2010 Lancieux et al.
 2010/0311028 A1 12/2010 Bell, III et al.
 2010/0316974 A1 12/2010 Yau et al.
 2011/0077635 A1 3/2011 Bonn

2011/0098697 A1 4/2011 Brannan
 2011/0104632 A1 5/2011 Colby
 2011/0130689 A1 6/2011 Cohen et al.
 2011/0137156 A1 6/2011 Razzaque et al.
 2011/0200960 A1 8/2011 Colby
 2011/0200961 A1 8/2011 Colby
 2011/0238060 A1 9/2011 Lee, Jr.
 2011/0238061 A1 9/2011 Van Der Weide
 2011/0244417 A1 10/2011 Hilsen et al.
 2012/0046536 A1 2/2012 Cheung et al.
 2012/0053577 A1 3/2012 Lee
 2012/0100500 A1 4/2012 Gao
 2012/0143180 A1 6/2012 Lee, Jr.
 2013/0017507 A1 1/2013 Moffson et al.
 2013/0150843 A1 6/2013 Berna et al.
 2013/0172731 A1 7/2013 Gole
 2013/0197355 A1 8/2013 Lee et al.
 2014/0046316 A1 2/2014 Ludtkow et al.
 2014/0066917 A1 3/2014 Cosman
 2014/0081260 A1 3/2014 Cosman
 2014/0093838 A1 4/2014 Carmichael et al.
 2014/0121658 A1 5/2014 Cosman
 2014/0186796 A1 7/2014 Suttin
 2014/0193771 A1 7/2014 Dolfi et al.
 2014/0257272 A1 9/2014 Clark, III et al.
 2014/0276033 A1 9/2014 Brannan et al.
 2014/0378964 A1 12/2014 Pearson
 2015/0018801 A1 1/2015 Lazarof
 2015/0018822 A1 1/2015 Racz
 2015/0038956 A1 2/2015 Amabile
 2015/0133909 A1 5/2015 Van Der Weide
 2015/0147714 A1 5/2015 Daon
 2015/0150631 A1 6/2015 Lee
 2015/0265371 A1 9/2015 Kim
 2015/0272671 A1 10/2015 Van Der Weide
 2015/0351831 A1 12/2015 Janssen
 2015/0374456 A1 12/2015 Colby
 2015/0374457 A1 12/2015 Colby
 2016/0058508 A1 3/2016 Brannan
 2016/0106518 A1 4/2016 Choi et al.
 2016/0151113 A1 6/2016 Kim et al.
 2016/0157967 A1 6/2016 Kim et al.
 2016/0324597 A1 11/2016 Colby
 2016/0367318 A1 12/2016 Van Der Weide
 2017/0014185 A1 1/2017 Lee, Jr.
 2017/0039489 A1 2/2017 Reh et al.
 2017/0231718 A1 8/2017 Wohrle
 2017/0290554 A1 10/2017 Merritt
 2017/0360528 A1 12/2017 Colby
 2018/0014910 A1 1/2018 Colby
 2018/0078309 A1 3/2018 Van Der Weide
 2018/0091169 A1 3/2018 Colby
 2018/0125579 A1 5/2018 Van Der Weide
 2018/0132934 A1 5/2018 Van Der Weide
 2018/0153640 A1 6/2018 Colby
 2018/0261922 A1 9/2018 Behdad et al.
 2018/0318038 A1 11/2018 Colby
 2019/0029751 A1 1/2019 Hancock et al.
 2019/0247144 A1 8/2019 Colby
 2020/0060761 A1 2/2020 Cao
 2020/0197089 A1 6/2020 Cao
 2021/0007829 A1 1/2021 Colby
 2022/0047356 A1 2/2022 Colby
 2022/0087744 A1 3/2022 Colby
 2023/0116948 A1 4/2023 Colby

FOREIGN PATENT DOCUMENTS

AU 2017/202885 7/2019
 AU 2019/204646 12/2020
 AU 2020217369 12/2021
 AU 2020287387 B2 2/2024
 CA 2761652 10/2019
 CA 2939815 8/2020
 CA 2939821 8/2020
 CN 107822710 3/2018
 DE 19510294 10/1996
 EP 1210022 4/2006
 EP 2301469 A1 3/2011

(56)

References Cited

FOREIGN PATENT DOCUMENTS

EP	2429444	3/2012
EP	3228272	10/2017
EP	3979938	4/2022
EP	2429444 B1	2/2024
FR	2925289	1/2011
GB	2057888	4/1981
MX	357449	7/2018
WO	2004084748 A1	10/2004
WO	WO 2006/031317	6/2006
WO	WO 2007/057902	5/2007
WO	2007057902 A3	7/2007
WO	WO 2007/085719	8/2007
WO	WO 2008/067334	6/2008
WO	WO 2008/128720	10/2008
WO	WO 2010/132368	11/2010
WO	2011017168 A2	2/2011
WO	WO 2014/143014	9/2014
WO	WO2014/152519	9/2014
WO	2017103209 A1	6/2017
WO	2020247885	12/2020
WO	2020247953	12/2020
WO	2022093177	5/2022

OTHER PUBLICATIONS

European Patent Office, European Search Report including the European Search Opinion and Supplementary European Search Report for Application No. EP 10775341.0, Jul. 11, 2014, 8 pages.
Blankenship et al., "Third Molar Development in the Estimation of Chronologic Age in American Blacks as Compared with Whites," *Journal of Forensic Science*, 2007, 52(2):428-33, Memphis, TN 38163, USA.

Gordon et al., "The effects of local hypothermia on odontogenesis," *Journal of Oral Surgery*, Apr. 1979, 37(4):235-44, PubMed—indexed for Medline, USA.

Silvestri et al., "Prevention of Third Molar Development in Dog with Long Pulse Diode Laser: A Preliminary Report," *Lasers Surg. Med.*, 2007, 39(8):674-7 (ISSN: 0196-8092), Boston, MA 02111, USA.

Silvestri et al., "Prevention of Third Molar Tooth Development in Neonate Rat with a Long Pulse Diode Laser," *Lasers Surg. Med.*, 2004, 35(5):385-91 (ISSN: 0196-8092), Boston, MA 02111, USA.
Qi H., "CT-guided microwave ablation through the lungs for treating liver tumors near the diaphragm," *Clinical Research Paper, Oncotarget Journal*, 2017, vol. 8, (No. 45), pp. 79270-79278, 9 pages, www.impactjournals.com/oncotarget/.

Patent Cooperation Treaty, "International Preliminary Report on Patentability" for PCT/US2020/036508, May 18, 2021, 7 pages.

Patent Cooperation Treaty, International Search Report and Written Opinion for PCT/US2020/036508, Nov. 17, 2020, 18 pages.

Patent Cooperation Treaty, International Search Report and Written Opinion for PCT/US2020/036705, Nov. 17, 2020, 18 pages.

Itero, "OrthoCAD, A Digitally Perfect Orthodontic Impression", in-house text transcript of previous NPL submission of same title, Jul. 24, 2018, 3 pages.

Patent Cooperation Treaty, "International Search Report" and "Written Opinion" for PCT/US2020/057383, Jul. 22, 2021, 9 pages.

Patent Cooperation Treaty, "International Search Report" for PCT/IL2006/001329, Mar. 7, 2007, 6 pages.

Pogrel, M.A et al.; "White paper on third molar data;" American Association of Oral and Maxillofacial Surgeons; Mar. 2007; http://www.aaoms.org/docs/third_molar_white_paper.pdf; pp. 1-25; USA.
AAOMS; "White paper on third molar data," Article, Mar. 2007, 25 pages. Published at www.aaoms.org.

AAOMS; "Advocacy white paper on evidence based third molar surgery;" American Association of Oral and Maxillofacial Surgeons; Nov. 10, 2011; USA.

Ahmad et al., "Caries Experience and Periodontal Pathology in Erupting Third Molars," *Journal of Oral and Maxillofacial Surgery*, 2008; 66(5):948-53 (ISSN: 1531-5053). Chapel Hill, NC 27599, USA.

Al-Khateeb et al., "Pathology Associated with Impacted Mandibular Third Molars in a Group of Jordanians," *Journal of Oral and Maxillofacial Surgery*, 2006; 64(11): 1598-602. Irbid, Jordan.

American Association of Oral and Maxillofacial Surgeons, "White Paper on Third Molar Data," Article, Mar. 2007, 25 pages. Published at www.aaoms.org.

American Dental Association, Abstract of "Growth in Dental Spending Expected to Slow in 2009," American Dental Association, Feb. 26, 2009, 1 page. Published at www.ada.org.

American Public Health Association, "Opposition to Prophylactic Removal of Third Molars (Wisdom Teeth)," Policy Statement Database, Policy No. 20085, Oct. 28, 2008. Washington, DC 20001, USA.

Askboots, "Removing Wisdom Teeth," Web Article, 9 page, Sep. 21, 2007. BMJ Publishing Group Limited. Published at www.askboots.com.

Australian Government; IP Australia; Patent Examination Report No. 1 (2010247874); Jan. 2, 2014; 3 pages.

Bataineh, A.B., Abstract of "Sensory Nerve Impairment Following Mandibular Third Molar Surgery," *Journal of Oral and Maxillofacial Surgery*, 2001; 59(9): 1012-7 (ISSN: 0278-2391). Irbid, Jordan.

Blakey et al., "Impact of Removal of Asymptomatic Third Molars on Periodontal Pathology," *Journal of Oral and Maxillofacial Surgery*, 2009; 67(2): 245-50. Chapel Hill, NC 27599, USA.

Blakey et al., "Periodontal Pathology Associated with Asymptomatic Third Molars" *Journal of Oral and Maxillofacial Surgery*, 2002; 60(11): 1227-33 (ISSN: 0278-2391). Chapel Hill, NC 27599, USA.

Blondeau et al., "Extraction of Impacted Mandibular Third Molars: Postoperative Complications and Their Risk Factors," *Journal of the Canadian Dental Association*, Jun. 2007, vol. 73, No. 4, (ISSN: 1488-2159).

Brace et al., "Pulmonary Thermal Ablation: Comparison of Radiofrequency and Microwave Devices by Using Gross Pathologic and CT findings in a Swine Model," *Radiology*, Jun. 2009, vol. 251, 705-11. Madison, WI 53792.

Brace, C.L., "Microwave Ablation Technology Avoids Problems that Plague RFA, Offers Promise for New Applications," *Diagnostic Imaging*, Mar. 13, 2006, 2 pages. Madison, WI 53792.

Bui et al., "Types, Frequencies, and Risk Factors for Complications After Third Molar Extraction," *Journal of Oral and Maxillofacial Surgery*, 2003; 61:1379-1389, Chapel Hill, NC 27599, USA.

Christiaens et al., "Complications After Third Molar Extractions: Retrospective Analysis of 1,212 Teeth," *Rev Stomatol Chir Maxillofac*, 2002; 103(5):269-74 (ISSN: 0035-1768). Belgique, France.

Dodson, T. B., "Response to APHA C3: Opposition to Prophylactic Removal of Third Molars," Article, Oct. 26, 2008, 3 pages, Massachusetts General Hospital. Boston, MA 02114, USA.

Dodson et al.; "Introduction;" © 2012 American Association of Oral and Maxillofacial Surgeons; © 2012 American Association of Oral and Maxillofacial Surgeons <http://dx.doi.org/10.1016/j.joms.2012.04.022>; *Journal of Oral and Maxillofacial Surgery*, vol. 70, No. 9, Supplement 1, Sep. 2012; pp. S2-S3; USA.

Dodson et al.; "Summary of the proceedings of the third molar multidisciplinary conference;" © 2012 American Association of Oral and Maxillofacial Surgeons; *Journal of Oral and Maxillofacial Surgery*, <http://dx.doi.org/10.1016/j.joms.2012.05.001>; vol. 70, No. 9, Supplement 1, Sep. 2012; pp. S66-S69; USA.

Eklund et al., "Third-Molar Removal Patterns in an Insured Population," *Journal of the American Dental Association*, vol. 132, No. 4, 469-475, 2001.

Elter et al., "Third Molars Associated with Periodontal Pathology in the Third National Health and Nutrition Examination Survey," *Journal of Oral and Maxillofacial Surgery*, 2004; 62(4):440-5. Chapel Hill, NC 27599, USA.

Elter et al., "Third Molars Associated with Periodontal Pathology in Older Americans," *Journal of Oral and Maxillofacial Surgery*, Feb. 2005; 63(2): 179-84. Chapel Hill, NC 27599, USA.

Fox, M.; "Shots kill budding wisdom teeth, study suggests;" © NBCNews.com; Apr. 2013; http://vitals.nbcnews.com/_news/2013/04/03/17584797-shots-kill-budding-wisdom-teeth-studysuggests%0D%0A%0D%0ANBCNews.com; New York, NY.

(56)

References Cited

OTHER PUBLICATIONS

- Friedman J.W., "Opposition to Prophylactic Removal of Third Molars," APHA Oral Health Section, Feb. 18, 2008, pp. 1-10.
- Friedman, J.W., "Containing the Cost of Third-Molar Extractions: A Dilemma for Health Insurance." *Public Health Reports*, vol. 98, No. 4, pp. 376-384, Jul.-Aug. 1983. Los Angeles, CA 90057, USA.
- Friedman, J.W., "The Prophylactic Extraction of Third Molars: A Public Health Hazard," *American Journal of Public Health*, vol. 97, No. 9, Sep. 2007. Los Angeles, CA 90064, USA.
- Gordon N.C. et al., "The effects of local hypothermia on odontogenesis," Abstract; *Journal of Oral Surgery*, Apr. 1979; 37(4):235-44; PubMed—indexed for Medline; USA.
- Guralnick W.C. et al., "Removal of third molars;" National Institutes of Health Consensus Development Conference Statement; Nov. 28-30, 1979; NIH Consensus Statement Online Nov. 28-30, 1979; 2(11):65-68; USA.
- Harrison, L.; "Nerve blocks in children may destroy future molars;" © 2013 WebMD, LLC; Medscape Medical News; Apr. 8, 2013; www.medscape.com; pp. 1-2, USA.
- Hicks, E.P., "Third Molar Management: A Case Against Routine Removal in Adolescent and Young Adult Orthodontic Patients," *Journal of Oral and Maxillofacial Surgery*, 1999; 57(7):831-6. Lexington, KY, USA.
- Huang et al., "Third-Molar Extraction as a Risk Factor for Temporomandibular Disorder," *Journal of the American Dental Association*, vol. 137, No. 11, pp. 1547-1554. American Dental Association, 2006.
- Itero, "OrthoCAD," "a digitally perfect orthodontic impression," www.cadentic.com, © 2009 Cadent, Ltd., 7 pages.
- Jiang, P., "Ten-Year Insurance Claims Data Study Reveals Increase in Utilization of Surgical Extraction Codes," *Online Periodical*, Winter 2006/2007, 2 pages, Delta Dental of Minnesota, Eagan, MN 55122, USA.
- Kajji et al., "Presence of Third Molar Germs in Orthodontic Patients in Japan," *American Journal of Orthodontics & Dentofacial Orthopedics*, vol. 119, Issue 3, pp. 245-250, Mar. 2001.
- Kaminishi et al., "A 10-year Comparative Study of the Incidence of Third Molar Removal in the Aging Population," *Journal of Oral Maxillofacial Surgery*, 2006; 64(2): 173-4 (ISSN: 0278-2391). Los Angeles, CA, USA.
- Kaminishi et al., "New Considerations in the Treatment of Compromised Third Molars," *Journal of the California Dental Association*, 2004; 32(10):823-5 (ISSN: 1043-2256). Los Angeles, CA.
- Kan et al., "Residual Periodontal Defects Distal to the Mandibular Second Molar 6-36 Months after Impacted Third Molar Extraction," *J. Clinical Periodontal*, Nov. 2002; 29(11):1004-11. Hong Kong SAR, China.
- Kiukkonen, "Toxicity of dioxin to developing teeth and salivary glands," Helsinki University Biomedical Dissertations No. 77, Jun. 16, 2006, 94 pages.
- Knutsson et al., "Dentists' Decision on Prophylactic Removal of Mandibular Third Molars: A 10-year Follow-Up Study," *Community Dental and Oral Epidemiology*, 2001; 29(4):308-14. Malmö University, Sweden.
- Kunkel et al., "Guideline: Surgical Removal of Third Molars," Agency for Quality in Dentistry, Apr. 2006, pp. 1-16. Cologne, Germany 50931.
- Kunkel et al., "Severe Third Molar Complications Including Death—Lessons from 100 Cases Requiring Hospitalization," *Journal of Oral Maxillofacial Surgery*, 2007; 65(9):1700-6. Mainz, Germany.
- Kunkel et al., "Third Molar Complications Requiring Hospitalization," *Oral Surg. Oral Med. Oral Pathol. Oral Radiol Endod.*, 2006; 102(3):300-6 (ISSN: 1528-395X). Mainz, Germany.
- Kusek, K.; "Researchers use light to coax stem cells to repair teeth;" © 2013 President and Fellows of Harvard College; Harvard School of Engineering and Applied Sciences; Wyss Institute; May 28, 2014; http://www.seas.harvard.edu/news/2014/05/researchers-use-light-to-coax-stem-cells-to-repair-teeth; pp. 1-6; Cambridge/Boston, MA, USA.
- Langsten et al., "The Impact of Retained Third Molars on the Deployed Airman," *Military Medicine*, vol. 173, Supplement 1, Jan. 2008, pp. 27-28(2). Publisher: AMSUS, Eglin Air Force Base, FL 32542.
- Levin, D.; "Nipping wisdom teeth in the bud: childhood anesthesia may thwart the development of third molars;" © 2014 Tufts University; Tufts Dental Medicine, Fall 2013; http://now.tufts.edu/articles/nipping-wisdom-teeth-bud; 1 page; Boston, MA, USA.
- McArdle et al., "Distal Cervical Caries in the Mandibular Second Molar: An Indication for the Prophylactic Removal of the Third Molar?" *British Journal of Oral Maxillofacial Surgery*, Feb. 2006; 44(1):42-5. London, UK.
- Moss et al., "Third Molar Periodontal Pathology and Caries in Senior Adults," *Journal of Oral Maxillofacial Surgery*, 2007; 65(1):103-8 (ISSN: 0278-2391). Chapel Hill, NC 27599, USA.
- Nance et al., "Change in Third Molar Angulation and Position in Young Adults and Follow-Up Periodontal Pathology," *Journal Oral Maxillofacial Surgery*, Mar. 2006; 64(3):424-8. Chapel Hill, NC 27599, USA.
- National Institutes of Health, "Removal of Third Molars," NIH Consensus Statement Online, Nov. 28-30, 1979; 2(11):65-8.
- Park et al., "Cortical Integrity of the Inferior Alveolar Canal as a Predictor of Paraesthesia After Third-Molar Extraction," *Journal of American Dental Association*, 2010; vol. 141, No. 3, 271-8.
- Patent Cooperation Treaty, "Notification of Transmittal of International Preliminary Report on Patentability," and "International Preliminary Report on Patentability," for PCT/US13/32357, at least as early as May 8, 2015, 14 pages.
- Patent Cooperation Treaty; International Searching Authority; International Search Report and Written Opinion of PCT/US13/32357; Jun. 5, 2013; 24 pages.
- Patent Cooperation Treaty, International Preliminary Report on Patentability, PCT/US10/34259, Aug. 31, 2011.
- Patent Cooperation Treaty, Written Opinion International Search Report, PCT/US10/34259, Sep. 14, 2010.
- Pogrel, M.A.; "What is the effect of timing of removal on the incidence and severity of complications?"; © 2012 American Association of Oral and Maxillofacial Surgeons; *Journal of Oral and Maxillofacial Surgery*; http://dx.doi.org/10.1016/j.joms.2012.04.028; vol. 70, No. 9, Supplement 1, Sep. 2012; pp. S37-S40; USA.
- Pratt et al., "Indications for Third Molar Surgery," *J. R. Coll. Surg. Edinb.*, vol. 43, Apr. 1998, pp. 105-108. Portsmouth, UK.
- Rakprakitkul, S., "Pathologic Changes in the Pericoronar Tissues of Unerupted Third Molars," *Quintessence International*. 2001; 32(8):633-8 (ISSN: 0033-6572). Rajthevee, Bangkok, Thailand 10400, 1 page.
- Ricketts, R.M. et al.; "Third molar enucleation: diagnosis and technique;" *Journal—California Dental Association*; Apr. 1976; http://www.ncbi.nlm.nih.gov/pubmed/1074850; pp. 52-57; USA.
- Sarnat et al., "Developmental Stages of the Third Molar in Israeli Children," *Pediatric Dentistry*, 2003; 25(4):373-7 (ISSN: 0164-1263). Tel Aviv, Israel.
- Shugars et al., "Incidence of Occlusal Dental Caries in Asymptomatic Third Molars" *Journal Oral Maxillofacial Surgery*, 2005; 63(3):341-6 (ISSN: 0278-2391). Chapel Hill, NC 27599, USA.
- Silvestri et al., "Selectively Preventing Development of Third Molars in Rats Using Electrosurgical Energy," *Journal of the American Dental Association*, vol. 135, No. 10, 1397-1405. Boston, MA 02111, USA, Oct. 2004.
- Silvestri et al., "The Unresolved Problem of the Third Molar: Would People be Better Off Without It?" *Journal of the American Dental Association*, Apr. 2003, vol. 134, No. 4, 450-5. Boston, MA 02111, USA.
- Song et al., "The Effectiveness and Cost-Effectiveness of Prophylactic Removal of Wisdom Teeth," Executive Summary, Health Technology Assessment, 2000; vol. 4: No. 15.
- Swée, J. et al.; "Inferior alveolar nerve block and third-molar agenesis: a retrospective clinical study;" © 2014 American Dental Association, all rights reserved; *JADA* 2013; 144(4); http://jada.ada.com; pp. 389-395; USA.
- U. S. Department of Labor, "Dentists," Webpage, 3 pages, last modified Dec. 17, 2009.

(56)

References Cited

OTHER PUBLICATIONS

University of York, the, "Prophylactic Removal of Impacted Third Molars: Is It Justified?" Effectiveness Matters, vol. 3, Issue 2, Oct. 1998. Heslington, York, UK.

Venta et al., "Change in Clinical Status of Third Molars in Adults during 12 Years of Observation," *Journal Oral Maxillofacial Surgery*, 1999; 57(4):386-9 (ISSN: 0278-2391). Helsinki, Finland.

Venta et al., "Malpractice Claims for Permanent Nerve Injuries Related to Third Molar Removals," *Acta Odontol. Scand.*, 1998; 56(4):193-6 (ISSN: 0001-6357). Helsinki, Finland.

Voegelin et al., "Complications during and after Surgical Removal of Mandibular Third Molars. Impact of Patient Related and Anatomical Factors," *Schweiz Monatsschr Zahnmed*, 2008; 118(3):192-8. Bern, Switzerland.

Wealthy Dentist, the, "Dental Marketing: Dentists Refer Some Wisdom Tooth Extractions to Oral Surgeons;" *The Wealthy Dentist.com*; at least as early as 2008; pp. 1-5. Tiburon, CA 94920, USA. Wikipedia, "Third Molar," Wikipedia, Webpage, 9 pages, last modified Apr. 19, 2011.

Wikipedia, "Tooth Development," Wikipedia, Webpage, 16 pages, last modified Apr. 18, 2011.

AAOMS Task Force; "Summary of the Third Molar Clinical Trials: Report of the AAOMS Task Force for Third Molar Summary;" AAOMS; *Journal of Oral and Maxillofacial Surgery*; 2012; 70:2238-2248; USA; 11 pages.

Food & Drug Administration (FDA); "Review of Published Literature Between 2008 and 2018 of Relevance to Radiofrequency Radiation and Cancer;" FDA website; Feb. 2020; 113 pages; <https://www.fda.gov/media/135043/download>.

Henry; "Enucleation of developing mandibular third molar by lateral trepanation;" Article; *Proc., Royal Society Medicine* 62:837-839; © 1969; Elsevier, United Kingdom; 3 pages; available online at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1811173/pdf/procrsmed00306-0105.pdf>.

Henry; "Prophylactic Enucleation of Wisdom Tooth Follicles;" Letter to the Editor; *Lancet* 227:921; © 1936; Elsevier, United Kingdom; 1 page.

Henry; "Prophylactic Odontectomy of the Developing Mandibular Third Molar: A New Operation;" Article; *American Journal of Orthodontics and Oral Surgery*; vol. 24, Iss. 1, Jan. 1938, pp. 72-84; first page submitted herewith; full 13 page article is available online as of Jun. 17, 2004, at: <https://www.sciencedirect.com/science/article/abs/pii/S009663473890033X?via%3Dihub>.

Hines-Peralta, et al.; "Microwave Ablation: Results with a 2.45-GHz Applicator in Ex Vivo Bovine and In Vivo Porcine Liver;" *Journal of Radiology* 239(1):94-102; Abstract; © 2006; 1 page, Oak Brook, Illinois, USA; available at: <https://pubmed.ncbi.nlm.nih.gov/16484351/>.

Hojjatollah, et al.; "Antenna Designs for Microwave Tissue Ablation;" Author manuscript; PubMed Central; Feb. 27, 2019; Department of Electrical and Computer Engineering, Kansas State University, Manhattan, Kansas, USA; 43 pages.

International Commission on Non-Ionizing Radiation Protection (ICNIRP); "RF EMFS 100kHz-300GHz;" ICNIRP Presentation; 2022; 3 pages; available at: <https://www.icnirp.org/en/frequencies/radiofrequency/index.html>KIM; "Understanding the Nuances of Microwave Ablation for More Accurate Post-Treatment Assessment;" *Future Oncology* 14(17):1755-1764; © 2018; 10 pages; available at: https://www.futuremedicine.com/doi/full/10.2217/fon-2017-0736?rfr_dat=cr_pub++0pubmed&url_ver=Z39.88-2003&rfr_id=ori%3Arid%3Acrsref.org.

Kim; "Understanding the Nuances of Microwave Ablation for More Accurate Post-Treatment Assessment;" *Future Oncology* 14(17):1755-1764; © 2018; 10 pages; available at: https://www.futuremedicine.com/doi/full/10.2217/fon-2017-0736?rfr_dat=cr_pub++0pubmed&url_ver=Z39.88-2003&rfr_id=ori%3Arid%3Acrsref.org.

Ling, et al.; "Which Procedure is Better: Germectomy or Surgical Removal of Mandibular Third Molar?"; *Int. J. of Oral Maxillofacial Surgery* 46:110; Abstract; 2017; 2 pages.

Lubner et al.; "Microwave Tumor Ablation: Mechanism of Action, Clinical Results, and Devices"; *Journal of Vascular and Interventional Radiology* 21(* Suppl.):S192-203; 2010; 28 pages; available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3065977/>.

Nanci; "Ten Cate's Oral Histology 7th Edition: Development, Structure, and Function;" Elsevier Mosby; © 2008; St., Louis, MO; 3 pages submitted herewith (including cover, publication page, and table of contents).

Shoshani-Dror, et al.; "Controversy regarding the need for prophylactic removal of impacted third molars: An overview;" Article; Quintessence Verlags-GmbH; © Sep. 2018; vol. 49, No. 8, 653-662; 10 pages; Online publication; document submitted is a personal PDF created for Leigh Colby, Account ID 2617823, created Sep. 2, 2022.

Simo, et al.; "Microwave Ablation Using 915-MHz and 2.45-GHz Systems: What Are the Differences?"; *HPB(Oxford)*; 15(12):991-996; 2013; 13 pages; available at: [https://www.hpbonline.org/article/S1365-182X\(15\)31352-6/fulltext](https://www.hpbonline.org/article/S1365-182X(15)31352-6/fulltext).

Vranckx, et al.; "Prophylactic vs. Symptomatic Third Molar Removal: Effects on Patient Postoperative Morbidity;" Feature Article; the *Journal of Evidence-Based Dental Practice*; © Sep. 2021; vol. 21, No. 3; 13 pages; Sage Publishing, Newbury Park, California; available online at: <https://www.sciencedirect.com/science/article/pii/S1532338221000579?via%3Dihub>.

Yang, Deshan; "Measurements, Antenna Design and Advanced Computer Modeling for Microwave Tissue Ablation"; Ph.D. Dissertation; University of Wisconsin—Madison; 2005; Madison, WI, USA; 272 pages.

Shugars et al.; "Assessment of Oral Health-Related Quality of Life Before and After Third Molar Surgery"; *Basic and Patient-Oriented Research*; 2006; pp. 1721-1730; vol. 64; American Association of Oral and Maxillofacial Surgeons; USA; 10 pages.

Sifuentes-Cervantes et al.; "Third molar surgery: Past, present, and the future"; *Oral Surgery Oral Medicine, Oral Pathology, and Oral Radiology*; 2021; pp. 523-531; vol. 132; Elsevier Inc.; 9 pages.

Sigron et al.; "The most common complications after wisdom-tooth removal"; *Research Science*; Oct. 2014; pp. 1041-1046; vol. 124; *Swiss Dental Journal*; Sweden; 6 pages.

Silvestri et al.; "Inferior alveolar nerve block and third-molar agenesis"; *JADA Continuing Education*; 2013; pp. 389-395; American Dental Association; USA; 7 pages.

Singh et al.; "Management of asymptomatic impacted wisdom teeth: a multicenter comparison"; *British Journal of Oral and Maxillofacial Surgery*; 1996; pp. 389-393; vol. 34; The British Association of Oral and Maxillofacial Surgeons; Scotland, UK; 5 pages.

Singh et al.; "The predictivity of mandibular third molar position as a risk indicator for pericoronitis: A prospective study"; *National Journal of Maxillofacial Surgery*; 2018; pp. 215-221; vol. 9; Wolters Kluwer—Medknow; 7 pages.

Slade et al.; "The Impact of Third Molar Symptoms, Pain, and Swelling on Oral Health-Related Quality of Life"; *Journal of Oral and Maxillofacial Surgery*; 2004; pp. 1118-1124; vol. 62; American Association of Oral and Maxillofacial Surgeons; USA; 7 pages.

Smith et al.; "Microbial Contamination and the Sterilization/Disinfection of Surgical Guides Used in the Placement of Endosteal Implants"; *The International Journal of Oral & Maxillofacial Implants*; 2011; pages unnumbered copyright warning page and 274-281; vol. 26, No. 2; Quintessence Publishing Co, Inc.; USA; 9 pages.

Snyder, et al.; "Pain Medication as an Indicator of Interference with Lifestyle and Oral Function During Recovery After Third Molar Surgery"; *Journal of Oral and Maxillofacial Surgery*; Aug. 2005; pp. 1130-1137; vol. 63, No. 8; © 2005 AAOMS; USA; 8 pages.

Staderini et al.; "How to Manage Impacted Third Molars: Germectomy or Delayed Removal? A Systemic Literature Review"; *Medicina*; 2019; pp. 1-14; vol. 55, No. 79; MDPI; Switzerland; 14 pages.

Steel et al.; "Current thinking in lower third molar surgery"; *British Journal of Oral and Maxillofacial Surgery*; 2021; available at <https://www.sciencedirect.com/science/article/abs/pii/S0266435621002667> from Jul. 30, 2021; 9 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Steel, et al.; "Current thinking in lower third molar surgery"; *British Journal of Oral and Maxillofacial Surgery*; 2022; pp. 257-265; vol. 60; © The British Association of Oral and Maxillofacial Surgeons; Elsevier Ltd.; 9 pages.
- Sun et al.; "The characteristics of adjacent anatomy of mandibular third molar germs: a CBCT study to assess the risk of extraction"; *Nature.com/scientificreports/*; Oct. 26, 2017; © Sun et al.; Nature; 7 pages.
- Svendson, et al.; "Third molar impaction—a consequence of late M3 mineralization and early physical maturity"; *European Journal of Orthodontics*; Feb. 1, 1988; vol. 10, No. 1; European Orthodontic Society; Sweden; 2 pages.
- Syed et al.; "Prevalence of Distal Caries in Mandibular Second Molar Due to Impacted Third Molar"; *Journal of Clinical and Diagnostic Research*; Mar. 2017; pp. 28-30; vol. 11, No. 3; jcdr.net; 3 pages.
- Synan et al.; "Management of Impacted Third Molars"; *Oral and Maxillofacial Surgery Clinics of North America*; 2020; pp. 519-559; vol. 32; © 2020 Elsevier Inc.; 41 pages.
- Tambuwalla et al.; "An Evaluation of Pathologic Changes in the Follicle of Impacted Mandibular Third Molars"; *Journal of International Oral Health*; 2015; pp. 58-62; vol. 7, No. 4; 5 pages.
- Tassoker et al.; "Is There a Possible Association between Skeletal Face Types and Third Molar Impaction? A Retrospective Radiographic Study"; *Medical Principles and Practice*; 2019; pp. 70-74; vol. 28; Karger Open Access; Switzerland; 5 pages.
- The Business Research Company; "Dental Services Market by Type (General Dentistry, Orthodontics; and Prosthodontics and Oral Surgery), by End Use Industry, by Country, By Competitor and Regional Analysis Global Forecast to 2022"; *Dental Services Market Report*; Aug. 2019; The Business Research Company; available online at: <https://www.thebusinessresearchcompany.com/report/dental-services-market>; 3 pages.
- Tolman et al.; "Continued evaluation of the effects of a sclerosing agent on odontogenesis"; *Oral Surgery Oral Medicine, Oral Pathology, and Oral Radiology Journal*; Jul. 1972; pp. 172-174; 6 pages.
- TriAgencies Brochure, "Zero3 TBA blocks third molar formation," 2023, 2 pages.
- TriAgencies; "Literature Survey on Third Molar Removal"; at least as early as Jul. 2, 2023; 15 pages.
- Tuovinen, et al.; "Is the third molar erupting at a younger age than before?"; *Acta Odontologica Scandinavica*; 2022; pp. 20-3209; vol. 80, No. 3; <https://doi.org/10.1080/00016357.2021.1985167>; Taylor & Francis; 8 pages.
- University of York, the, "Prophylactic Removal of Impacted Third Molars: Is It Justified?" *Effectiveness Matters*, vol. 3, Issue 2, Heslington, York, UK, Oct. 1998, 4 pages.
- Vandeplass, et al.; "Does Retaining Third Molars Result in the Development of Pathology Over Time? A Systematic Review"; 2020; *Journal of Oral and Maxillofacial Surgery*; vol. 78; pp. 1892-1908; AAOMS; 17 pages.
- Venta et al.; "A device to predict lower third molar eruption"; *Oral Surgery, Oral Medicine, Oral Pathology*; 1997; pp. 598-603; vol. 84, No. 6; Mosby—Year Book, Inc.; USA; 6 pages.
- Venta et al.; "Clinical Outcome of Third Molars in Adults Followed During 18 Years"; *Journal of Oral and Maxillofacial Surgery*; 2004; pp. 182-185; vol. 62; American Association of Oral and Maxillofacial Surgeons; USA; 4 pages.
- Venta, Irja; "How Often Do Asymptomatic, Disease-Free Third Molars Need to Be Removed?"; *Journal of Oral and Maxillofacial Surgery*; ; 2012; pp. S41-S47; Elsevier Inc.; 7 pages.
- Versaci, Mary Beth; "Dental spending exceeds pre-pandemic levels in 2021"; *ADA News*; Jan. 9, 2023; USA; 2 pages.
- Vranckx et al.; "Artificial Intelligence (AI)-Driven Molar Angulation Measurements to Predict Third Molar Eruption on Panoramic Radiographs"; *International Journal of Environmental Research and Public Health*; May 25, 2020; pp. 1-13; vol. 17; www.mdpi.com/journal/ijerph; MDPI; Basel, Switzerland; 13 pages.
- Vranckx et al.; "Radiographic prediction of mandibular third molar eruption and mandibular canal involvement based on angulation"; *Orthodontics and Craniofacial Research*; 2019; pp. 118-123; vol. 22; John Wiley & Sons Ltd.; available online at: wileyonlinelibrary.com/ocr; 6 pages.
- Vranckx, Myrthel, et al. "Surgical experience and patient morbidity after third molar removal." *Journal of Stomatology, Oral and Maxillofacial Surgery* 123.3 (2022): 297-302.
- Vujicic, et al.; "State of the Dental Market: Outlook 2018"; ADA and HPI; 2017; pp. 1-50; © ADA; USA; 50 pages.
- Vujicic, et al.; "Value-based care in dentistry"; *Journal of American Dental Association*; Guest Editorial; Jun. 2023; pp. 449-452; vol. 154, No. 6; available at: <http://jada.ada.org>; ADA; USA; 3 pages.
- Waite et al.; "Surgical Management of Impacted Third Molars"; *Seminars in Orthodontics*; 1998; pp. 113-123; vol. 4, No. 2; W.B. Saunders Company; USA; 11 pages.
- Wealthy Dentist, the, "Dental Marketing: Dentists Refer Some Wisdom Tooth Extractions to Oral Surgeons," *The Wealthy Dentist.com*, at least as early as 2008, Tiburon, CA 94920, USA, 5 pages.
- White et al.; "Recovery After Third Molar Surgery: Clinical and Health-Related Quality of Life Outcomes"; *Journal of Oral and Maxillofacial Surgery*; 2003; pp. 535-544; vol. 61; American Association of Oral and Maxillofacial Surgeons; USA; 10 pages.
- White et al.; "Special Contribution: Third Molar Clinical Trials Annotated Bibliography"; *Journal of Oral and Maxillofacial Surgery*; 2016; pp. 4-12; vol. 74; American Association of Oral and Maxillofacial Surgeons; USA; 9 pages.
- White, Raymond P.; "Recovery after third-molar surgery"; *Advances in Orthodontics & Dentofacial Surgery*; 2004; p. 289; American Association of Orthodontists; USA; 1 page.
- Wikipedia; "Near and far field"; Wikipedia entry; page last edited on Sep. 8, 2018; published online at https://en.wikipedia.org/wiki/Near_and_far_field; 11 pages.
- Wise et al.; "Cellular, molecular, and genetic determinants of tooth eruption"; *Critical Reviews in Oral Biology & Medicine*; 2002; abstract of article on pp. 323-334; vol. 13, Issue 4; USA; 1 page.
- Wu, R.-X.; Tian, B.-M.; Gao, R.; Chen, F.-M.; "Non-Impacted Third Molars: Angels or Devils?"; *Journal of Clinical Medicine*; 2023; 12;4455. <https://doi.org/10.3390/jcm12134455>; 3 pages.
- Xiang et al.; "Impact of platelet-rich fibrin on mandibular third molar surgery recovery: a systemic review and meta-analysis"; *BMC Oral Health*; 2019; pp. 1-10; vol. 19, Issue 163; Springer Nature; available online at: <https://doi.org/10.1186/s12903-019-0824-3>; 10 pages.
- Yildirim et al.; "Pathologic changes in soft tissues associated with asymptomatic impacted third molars"; *OOOOE*; Jul. 2008; pp. 14-18; vol. 106, No. 1; Mosby Inc.; USA; 5 pages.
- Zandi et al.; "Evaluation of third molar development and its relation to chronological age: a panoramic radiographic study"; *Journal of Oral and Maxillofacial Surgery*; 2015 (published online Nov. 21, 2014); pp. 183-189; vol. 19; Springer-Verlag; Berlin/Heidelberg, Germany; 7 pages.
- Zhang et al.; "Articaine Infiltration Versus Lidocaine Inferior Alveolar Nerve Block for Anesthetizing Primary Mandibular Molars: A Randomized, Controlled, Double-Blind Pilot Study"; *Pediatric Dentistry*; Sep.-Oct. 2021; pp. 344-348; vol. 43, No. 5; American Academy of Pediatric Dentistry; USA; 5 pages.
- Zhang et al.; "Early Extraction: a silver bullet to avoid nerve injury in lower third molar removal"; *International Journal of Oral & Maxillofacial Surgery*; 2012; pp. 1280-1283; vol. 41; Elsevier Ltd.; available online at <http://www.sciencedirect.com>; 4 pages.
- Zhou-Xi et al; "Pathologies associated with the mandibular third molar impaction"; *Science Progress*; 2021; pp. 1-10; vol. 104, No. 2; © Zhou-Xi, et al.; Sage; 10 pages.
- ZZQ (Agency for Quality in Dentistry—a unit of the Institute of German Dentists); "Surgical removal of third molars"; Guideline; Apr. 2006; pp. 1-16; German Dental Association and National Association of Statutory Health Insurance Dentists; Germany; 16 pages.
- Chaffin et al.; "Review of Current U.S. Army Dental Emergency Rates"; *Military Medicine*; 2008; pp. 23-26; vol. 173, January Supplement 2008; Association of Military Surgeons of the U.S.; USA; 4 pages.

(56)

References Cited

OTHER PUBLICATIONS

Chaudhry et al.; Efficacy of adjuvant ozone therapy in reducing postsurgical complications following impacted mandibular third-molar surgery'; Oct. 2021; pp. 842-854; vol. 152, No. 10; <http://jada.ada.org>; Canada; 13 pages.

Cheifetz et al.; "Preface-Proceedings of the Third Molar Multidisciplinary Conference"; Article Preface; Sep. 2012; p. S1; vol. 70, No. 9, Supp. 1; Journal of Oral and Maxillofacial Surgery; vol. 70, No. 9, Supplement 1; American Association of Oral and Maxillofacial Surgeons; USA; 1 page.

Chen, et al.; "Revisit incidence of complications after impacted mandibular third molar extraction: A nationwide population-based cohort study"; PLOS One article; Feb. 22, 2021; pp. 1-13; available at: <https://doi.org/10.1371/journal.pone.0246625>; 13 pages.

Chiapasco et al.; "Germectomy or Delayed Removal of Mandibular Impacted Third Molars: The Relationship Between Age and Incidence of Complications"; Journal of Oral and Maxillofacial Surgeons; 1995; pp. 418-422; vol. 53; American Association of Oral and Maxillofacial Surgeons; USA; 5 pages.

Cho et al.; "Postoperative interventions to reduce inflammatory complications after third molar surgery: review of the current evidence"; Australian Dental Journal; May 12, 2017; pp. 412-419; vol. 62, No. 4; <https://doi.org/10.1111/adj.12526>; Australia; 16 pages.

Chossegros et al.; "Is lingual nerve protection necessary for lower third molar germectomy? A prospective study of 300 procedures"; International Journal of Maxillofacial Surgeons; 2002; pp. 620-624; vol. 31; © International Journal of Maxillofacial Surgeons; published by Elsevier Science Ltd.; USA; 5 pages.

Chuang et al.; "Risk Factors for Inflammatory Complications Following Third Molar Surgery in Adults"; Journal of Oral and Maxillofacial Surgery; 2008; pp. 2213-2218; vol. 66; American Association of Oral and Maxillofacial Surgeons; USA; 6 pages.

Cohen et al.; "Patients' Restrospective Preference for Extraction of Asymptomatic Third Molars"; Community Dentistry and Oral Epidemiology; 1990; pp. 260-263; vol. 18; DTIC File Copy of the Naval Dental Research Institute; USA; 8 pages.

Colby & Watson; "Fully Guided Tooth Bud Ablation in Pigs Results in Complete Tooth Bud Removal and Molar Agenesis"; Journal of Oral and Maxillofacial Surgery; 2022; pp. 456-466; vol. 81; American Association of Oral and Maxillofacial Surgeons; USA; 11 pages.

Colby, Leigh E.; "A New Paradigm for Third Molar Management. The Clinical Rationale Behind Fully Guided Third Molar Tooth Bud Ablation"; Dentistry Today; Sep. 11, 2023; p. 42.; 1 page.

Colby, Leigh E.; "Fully Guided Third Molar Tooth Bud Ablation in Pigs"; Journal of Oral and Maxillofacial Surgery; 2022; pp. 1522-1533; vol. 80; American Association of Oral and Maxillofacial Surgeons; USA; 12 pages.

Combes et al.; "Third molar-related morbidity in deployed Service personnel"; British Dental Journal; 2010; pp. 1-4; vol. 209, E6; British Dental Journal; Britain; 4 pages.

Cooper-Newland, Deborah L.; "Management of Impacted Third Molar Teeth"; The Journal of the Greater Houston Dental Society; Mar. 1996; pp. 10-12; vol. 67, No. 8; USA; 4 pages.

Cortigiano, Cameron; "Number of dentists by specialty type"; Becker's Dental + DSO Review; at least as early as Feb. 26, 2023; Becker's Dental + DSO website; 1 page.

Costich, Emmett R.; "Removal of third molars as a phase of preventive dentistry"; Dental Abstracts; Jan. 1968; pp. 9-21; vol. 13, No. 1; USA; 14 pages.

Cunha-Cruz et al.; "Recommendations for Third Molar Removal: A Practice-Based Cohort Study"; American Journal of Public Health; Apr. 2014; pp. 735-743; vol. 104, No. 4; American Journal of Public Health; USA; 9 pages.

D'Angeli, et al.; "The Evaluation of Further Complications after the Extraction of the Third Molar Germ: A Pilot Study in Paediatric Dentistry"; Healthcare; 2021; pp. 1-12; vol. 9; <https://doi.org/10.3390/healthcare9020121>; MDPI; 12 pages.

Da Costa et al.; "Is there justification for prophylactic extraction of third molars? A systematic review"; Braz Oral Res.; Mar.-Apr. 2013; pp. 183-188; vol. 27, No. 2; Sao Paulo, Brazil; 6 pages.

Dai et al.; 'Mapping the amelogenin protein expression during porcine molar crown development; Annals of Anatomy; pp. 1-7; vol. 234; Elsevier GmbH; 7 pages.

De Souza Povoia et al.; "Does the Coronectomy a Feasible and Safe Procedure to Avoid the Inferior Alveolar Nerve Injury during Third Molars Extractions? A Systematic Review"; Healthcare; 2021; pp. 1-13; vol. 9.; <https://doi.org/10.3390/healthcare9060750>; MDPI; Basel, Switzerland; 13 pages.

Deliverska et al.; "Complications After Extraction of Impacted Third Molars—Literature Review"; Journal of IMAB; 2016; pp. 1202-1211; vol. 22, Issue 3; Journal of IMAB; available online at: <http://www.journal-imab-bg.org>; 10 pages.

Dhongde et al.; "Assessment of Growth Status: Nolla's Dental Age v/s Chronological Age"; International Journal of Oral Health and Medical Research; Jan.-Feb. 2017; pp. 15-17; vol. 3, Issue 5; India; 3 pages.

Dodson et al.; "Impacted wisdom teeth"; Clinical Evidence; Aug. 29, 2014; pp. 1-15; vol. 2014, No. 1302; BMJ Publishing Group Ltd; 15 pages.

Dodson Thomas B.; "Surveillance as a Management Strategy for Retained Third Molars: Is It Desirable?"; Journal of Oral and Maxillofacial Surgery; ; 2012; pp. 20-24; vol. 70, Suppl. 1; American Association of Oral and Maxillofacial Surgeons; USA; 5 pages.

Dodson, Thomas B.; "How Many Patients Have Third Molars and How Many Have One or More Asymptomatic, Disease-Free Third Molars?"; Journal of Oral and Maxillofacial Surgery; 2012; pp. S4-S7; vol. 70; American Association of Oral and Maxillofacial Surgeons; USA; 4 pages.

Dodson, Thomas B.; "The Management of the Asymptomatic, Disease-Free Wisdom Tooth: Removal Versus Retention"; Atlas of Oral and Maxillofacial Surgery Clinics North America; 2012; pp. 169-176; vol. 20; Elsevier Inc.; 8 pages.

Dodson, Thomas B.; "Writing a Scientific Paper Is Not Rocket Science!"; Journal of Oral and Maxillofacial Surgeons; 2015; pp. S160-S169; American Association of Oral and Maxillofacial Surgeons; USA; 10 pages.

Donnell, et al.; "Mandibular third molars: 'naughty' or Nice?"; British Dental Journal; Apr. 10, 2020; pp. 506-507; vol. 228, No. 7; British Dental Association; UK; 2 pages.

Dunn et al.; "Dental Emergency Rates at Two Expeditionary Medical Support Facilities Supporting Operations Enduring and Iraqi Freedom"; Military Medicine; Jul. 2004; pp. 510-514; vol. 169, No. 7; Association of Military Surgeons; USA; 5 pages.

Edwards et al.; "Impact of third molar removal on demands for postoperative care and job disruption: does anaesthetic choice make a difference"; Annals of the Royal College of Surgeons; 1999; pp. 119-123; England; 5 pages.

Eklund, Stephen A.; "Trends in dental treatment, 1992-2007"; Journal of American Dental Association Continuing Education; Apr. 2010; pp. 391-399; vol. 141; American Dental Association (ADA); USA; 9 pages.

Ethicon; "Neuwave™ Microwave Ablation System—Build your ablation program with the only system that has it all"; Brochure No. 068741-170308; Ethicon; 2017; 2 pages.

Ethicon; "Neuwave™ Microwave Ablation System"; Product description; Ethicon; last updated Dec. 15, 2017; published online at: <https://www.ethicon.com/na/products/microwave-ablation/microwave-ablation/neuwave-microwave-ablation-system>; 5 pages.

Extended European Search Report issued in App. No. EP20818571; dated May 31, 2023; 5 pages.

Fernandes et al.; "Actuarial life-table analysis of lower impacted wisdom teeth in general dental practice"; Community Dentistry and Oral Epidemiol; 2009; pp. 58-67; vol. 38; John Wiley & Sons A/S; 10 pages.

Fernandes et al.; "Incidence of symptoms in previously symptom-free impacted lower third molars assessed in general dental practice"; BDJ; Sep. 4, 2009; pp. 1-21; vol. 207, E.10; Britain; available online at <https://doi.org/10.1038/sj.bdj.2009.804>; 21 pages.

Flick; "The Third Molar Controversy: Framing the Controversy as a Public Health Policy Issue"; Journal of Oral and Maxillofacial

(56)

References Cited

OTHER PUBLICATIONS

- Surgeons; 1999; pp. 438-444; vol. 57; American Association of Oral and Maxillofacial Surgeons; USA; 7 pages.
- Frenzel, Lou; "What's the Difference Between EM Near Field and Far Field?"; Article; Electronic Design; Jun. 6, 2012; published online at: <http://electronicdesign.com/energy/whatsdifferencebetweenemnearfieldandfarfield>; 6 pages.
- Friedman, Harold; "The Impacted Mandibular Third Molar Problem"; The Dental Students' Magazine; Dec. 1946; pp. 13-16, and 34; vol. 25 1946/47;; USA; 6 pages.
- Friedman, Jay W.; "When Abstinence is Evidence-Based"; 2012 National Oral Health Conference presentation; May 2, 2012; pp. 1-49; USA; 49 pages.
- Garaas et al.; "Prevalence of Visible Third Molars With Caries Experience or Periodontal Pathology in Middle-Aged and Older Americans"; Journal of Oral and Maxillofacial Surgeons; 2011; pp. 463-470; vol. 69; © American Association of Oral and Maxillofacial Surgeons; USA; 8 pages.
- Garvin, Jennifer; "Oral Surgeons are 'consistently' highest earning specialists"; ADA News; Mar. 6, 2017; p. 8; ADA News; USA; 1 page.
- Ghaemina H.; "Surgical removal versus retention for the management of asymptomatic disease-free impacted wisdom teeth (Review)"; Cochrane Database of Systematic Reviews; 2020; pp. 1-27; Iss. 5, Art. No. CD003879; DOI 10.102/14651858.CD003879.pub5; cochranelibrary.com; John Wiley & Sons, Ltd.; 27 pages.
- Goldberg et al.; "Complications after mandibular third molar surgery: a statistical analysis of 500 consecutive procedures in private practice"; Journal of American Dental Association (JADA)—Brief Reports; 1985; pp. 277-279; vol. 111; USA; 3 pages.
- Goldberg et al.; "Impacted Third Molar Referral Patterns"; JADA—Brief Reports; Sep. 1983; pp. 439-441; vol. 107; USA; 3 pages.
- Guo et al.; "The influence of impaction to the third molar mineralization in northwestern Chinese population"; Int J Legal Med. article; Feb. 16, 2014; pp. 659-665; vol. 128; Springer; 7 pages.
- Gupta et al.; "Recent Trends in the Market for Oral Surgeons, Endodontists, Orthodontists, Periodontists, and Pediatric Dentists"; ADA & HPI; Feb. 2017; pp. 1-14; ADA; USA; 14 pages.
- Gupta et al.; "Trends in Fees and Reimbursement Rates for the Most Common Procedures in Endodontics, Periodontics, and Oral Surgery"; ADA & HPI; Apr. 2017 (Revised); pp. 1-8; ADA; USA; 8 pages.
- Haddad et al.; "The Importance of Recognizing Pathology Associated with Retained Third Molars"; JCDA; Feb. 2006; pp. 41-45; vol. 72, No. 1; Canada; 5 pages.
- AAOMS; "Management of Third Molar Teeth"; AAOMS White Paper; 2016; p. 1-2; AAOMS; USA; 2 pages.
- AAOMS; "Parameters of Care: AAOMS Clinical Practice Guidelines for Oral and Maxillofacial Surgery (AAOMS ParCare) Sixth Edition"; Supplement to the Journal of Oral and Maxillofacial Surgery, Sixth Edition; Aug. 2017; cover and pp. e61-e64; vol. 75, No. 8, Suppl. 1; AAOMS; www.aaoms.org; USA; 5 pages.
- AAOMS; "Supporting Information to the Management of Patients with Third Molar Teeth"; AAOMS; 2016; pp. 1-3; AAOMS; Rosemont, IL, USA; 3 pages.
- AAOMS; "The Management of Impacted Third Molar Teeth," AAOMS Clinical Paper; 2017; p. 1-4; AAOMS; USA; 4 pages.
- AAOMS; "Third Molar Clinical Studies—Summary of Data"; AAOMS—Third Molar Multidisciplinary Conference; 2012; 2 pages; AAOMS; USA; 2 pages.
- AAOMS; "Wisdom Teeth Management"; Oral and maxillofacial surgeons: The experts in face, mouth and jaw surgery © 2005-2013; pp. 1-3; AAOMS E-book; USA; available online at: https://www.aaoms.org/images/uploads/pdfs/Ebook_Wisdom_Teeth_R.pdf; 3 pages.
- AAPD; "Guideline on Pediatric Oral Surgery"; American Academy of Pediatric Dentistry (AAPD) Reference Manual Clinical Guidelines; Revised 2010; pp. 238-245; V. 32, No. 6; AAPD; Chicago, IL, USA; 8 pages.
- AAPD; "Periodicity of Examination, Preventive Dental Services, Anticipatory Guidance/Counseling, and Oral Treatment for Infants, Children, and Adolescents"; The Reference Manual of Pediatric Dentistry; 2021; pp. 241-251; Chicago, Illinois, USA; 11 pages.
- AAPD; "The Reference Manual of Pediatric Dentistry: Definitions, Oral health policies, Recommendations, Endorsements, Resources"; 2021-2022; cover and pp. 245-246; © America's Pediatric Dentists—the Big Authority on little teeth®; USA; 3 pages.
- Abdullahi et al.; "Animal Models in Burn Research"; National Institutes of Health (NIH) Public Access; Author Manuscript; 2014; pp. 1-26; (available in PubMed Central on Sep. 1, 2015); USA; 26 pages.
- Abrahams, et al.; "Pediatric Odontogenic Tumors"; Oral and Maxillofacial Surgery Clinics of North America; 2016; copyright notice and pp. 45-58 of journal; vol. 28; © Elsevier Inc.; 15 pages.
- ADA & HPI; "Dental Fees, Results from the 2018 Survey of Dental Fees"; ADA, HPI, 2018 Survey of Dental Fees; © 2018 ADA; pp. 1-141; ADA; USA; 141 pages.
- ADA & HPI; "U.S. Dental Expenditures, 2017 Update"; ADA & HPI 2017 Update; © 2017 ADA; pp. 1-10; ADA; USA; 10 pages.
- ADA & HPI; "U.S. Dental Spending Down in 2020"; Health Policy Institute—infographic; accessed at least as early as Feb. 9, 2022; available online at: ada.org/hpi; USA; 1 page.
- ADA & HPI; "U.S. Dental Spending Up in 2021"; Health Policy Institute—infographic; accessed at least as early as Jun. 27, 2023; available online at: <https://www.ada.org/resources/research/health-policy-institute/us-dental-spending-in-2021>; USA; 1 page.
- Adeyemo; "Do pathologies associated with impacted lower third molars justify prophylactic removal? A critical review of the literature"; Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology; 2006; pp. 448-452; © 2006 Mosby, Inc.; 5 pages.
- Adeyomo, W. L.; "Indications for extraction of third molars: a review of 1763 cases"; Nigeria Postgrad Medical Journal; Mar. 15, 2008; abstract of article on pp. 42-46; vol. 1; Nigeria; 1 page.
- Akinbami et al.; "Dry Socket: Incidence, Clinical Features, and Predisposing Factors"; International Journal of Dentistry; 2014; pp. 1-7; vol. 2014; Article ID 796102; Hindawi Publishing Corporation; 7 pages.
- Al-Asfour; "Postoperative Infection after Surgical Removal of Impacted Mandibular Third Molars: An Analysis of 110 Consecutive Procedures"; Medical Principles and Practice; 2009; pp. 48-52; vol. 18; © 2008 Karger; Basel, Switzerland; available online at: www.karger.com/mpp; 5 pages.
- Al-Khateeb et al.; "Pathology associated with impacted mandibular third molars in a group of Jordanians"; J Oral Maxillofacial Surgery; Nov. 2006; 1598-1602; vol. 64, No. 11; USA; 5 pages.
- Allen, Bryana; "What to Know About Wisdom Teeth"; Delta Dental blog article; Mar. 8, 2018; Delta Dental of Washington; USA; 5 pages.
- American Academy of Pediatric Dentistry (AAPD); "Guideline on Pediatric Oral Surgery"; American Academy of Pediatric Dentistry (AAPD) Reference Manual Clinical Guidelines; Adopted 2005; cover pages with copyright notices and pp. 218-224 of reference manual; vol. 31, No. 6; AAPD; Chicago, IL, USA; 9 pages.
- American Association of Oral and Maxillofacial Surgeons (AAOMS); "Advocacy White Paper on Evidence Based Third Molar Surgery"; AAOMS; Nov. 10, 2011; AAOMS; USA; 6 pages.
- American Dental Association (ADA) & Health Policy Institute (HPI); "Dental Fees, Results from the 2016 Survey of Dental Fees"; ADA, HPI 2016 Survey of Dental Fees; © 2016 ADA; pp. 1-146; ADA; USA; 146 pages.
- Ash et al.; "A Study of Periodontal Hazards of Third Molars"; The Journal of Periodontology; Jul. 1962; pp. 209-219; vol. 33, Issue 3; USA; 11 pages.
- Assael, Leon A.; "Indications for Elective Therapeutic Third Molar Removal: The Evidence Is In"; Journal of Oral and Maxillofacial Surgery editorial; 2005; pp. 1691-1692; vol. 63; AAOMS; USA; 2 pages.
- Australian Examination Report No. 1 issued in App. No. AU2020475251; dated Jun. 29, 2023; 4 pages.
- Australian Examination Report No. 1 issued in App. No. AU2021282404; dated Feb. 10, 2023; 2 pages.
- Australian Examination Report No. 2 issued in App. No. AU2020475251; dated Sep. 14, 2023; 3 pages.

(56)

References Cited

OTHER PUBLICATIONS

Australian Notice of Acceptance issued in App. No. AU2021282404; dated Sep. 21, 2023; 3 pages.

Azab, et al.; Efficacy of secondary vs primary closure techniques for the prevention of postoperative complications after impacted mandibular third molar extractions; *JADA*; Oct. 2022; pp. 943-956.e48; vol. 153, No. 10; ADA; USA; 62 pages.

Baensch et al.; "Third Molar Complications in the Elderly—A Matched-Pairs Analysis"; *Journal of Oral and Maxillofacial Surgery*; I; 2017; pp. 680-686; vol. 75; © 2016 American Association of Oral and Maxillofacial Surgeons; 7 pages.

Bagheri et al.; "Extraction Versus Nonextraction Management of Third Molars"; *Oral and Maxillofacial Surgery Clinics of North America*; 2007; pp. 15-21; vol. 19; Elsevier Inc.; 7 pages.

Bailey et al.; Surgical techniques for the removal of mandibular wisdom teeth (Review); *Cochrane Database of Systematic Reviews*; 2020; pp. 1-164; www.cochranelibrary.com; John Wiley & Sons, Ltd.; 164 pages.

Barka et al.; "Radiographic evaluation of third molar genesis in Greek orthodontic patients"; *International Journal of General Medicine*; 2013; pp. 747-755; vol. 6; Dove Press; 9 pages.

Behbehani et al.; "Prediction of mandibular third-molar impaction in adolescent orthodontic patients"; *American Journal of Orthodontics and Dentofacial Orthopedics*; 2006; pp. 47-55; vol. 130, No. 1; © 2006 by the American Association of Orthodontists; USA; 9 pages.

Bhuyan, et al.; "Insight into the molecular pathogenesis of odontogenic lesions"; *Journal of Oral Biosciences*; 2021; pp. 35-44; vol. 63; © Japanese Association for Oral Biology; Published by Elsevier RV; 10 pages.

Boffano et al.; "Lingual nerve deficit following mandibular third molar removal: Review of the literature and medicolegal considerations"; *Journal of Oral and Maxillofacial Surgery*; vol. 113; No. 3; Mar. 2012; e10-e18; 9 pages.

Bonetti et al.; "Orthodontic Extraction: Riskless Extraction of Impacted Lower Third Molars Close to the Mandibular Canal"; *Journal of Oral and Maxillofacial Surgeons*; 2007; pp. 2580-2586; vol. 65; AAOMS; USA; 7 pages.

Bouloux et al.; "What is the Risk of Future Extraction of Asymptomatic Third Molars? A Systematic Review"; *Journal of Oral and Maxillofacial Surgery*; 2015; pp. 806-811; vol. 73; AAOMS; USA; 6 pages.

Brace, C. L.; "Microwave Ablation With a Triaxial Antenna: Results in ex vivo Bovine Liver"; Author Manuscript of Study Results; National Institute of Health; IEEE Trans Microw Theory Tech. Jan. 2005; 53(1):215-220; University of Wisconsin—Madison, Madison, WI, 53706 USA; 21 pages.

Bruce et al.; "An observational analysis of risk factors associated with symptomatic third molar teeth [version 2; peer review: 2 approved]"; *Wellcome Open Research* 2023; as published Jul. 17, 2023; p. 1-14; 14 pages.

Bruce et al.; "Age of patients and morbidity associated with mandibular third molar surgery"; *JADA*; Aug. 1980; pp. 240-245; vol. 101; USA; 6 pages.

Cahill et al.; "Tooth eruption: evidence for the central role of the dental follicle"; Abstract; *Journal of Oral Pathology*; Jul. 1980; abstract of article that was published on pp. 189-200; vol. 9, No. 4; Medline; 1 page.

Campbell, John H.; "Pathology Associated with the Third Molar"; *Oral and Maxillofacial Surgery Clinics of North America*; 2013; vol. 25; pp. 1-10; Elsevier Inc.; 10 pages.

Carter et al.; "Predictors of Third Molar Impaction: A Systematic Review and Meta-analysis"; *Journal of Dental Research*; 2016; pp. 267-276; vol. 95, No. 3; © International & American Association for Dental Research 2015; Sage Publications Inc.; 10 pages.

CenExcel JBR; "Wisdom Teeth Removal by the Numbers"; CenExcel clinical trial data; Mar. 13, 2015; available online at: <https://cenexcelresearch.com/jbr/wisdom-teeth-removal-numbers/>; Salt Lake City, UT, USA; 3 pages.

Centers for Medicare & Medicaid Services (CMS); "National Health Expenditures 2017 Highlights"; CMS.gov; 2018; pp. 1-3; Department of Health and Human Services (HHS); USA; 3 pages.

Centers for Medicare & Medicaid Services (CMS); "National Health Expenditures 2018 Highlights"; NHE; 2019; pp. 1-3; Department of Health and Human Services (HHS); USA; 3 pages.

Cervera-Espert et al.; "Coronectomy of impacted mandibular third molars: A meta-analysis and systematic review of the literature"; *Medicina Oral Patologia Oral y Circugia Bucal*; Jul. 1, 2016; pp. e505-e513; vol. 21, No. 4; Spain; 9 pages.

Harbaugh et al.; "Persistent Opioid Use After Wisdom Tooth Extraction"; *JAMA*; Aug. 7, 2018; pp. 504-506; vol. 320, No. 5; American Medical Association; USA; 3 pages.

Hill et al.; "Conservative, non-surgical management of patients presenting with impacted lower third molars: A 5-year study"; *British Journal of Oral and Maxillofacial Surgery*; 2006; pp. 347-350; vol. 44; Elsevier Ltd.; 4 pages.

Hounsborne et al.; "Prophylactic removal of impacted mandibular third molars: a systematic review and economic evaluation"; *National Institute for Health Research—Health Technology Assessment*; Jun. 2020; 146 pages; vol. 24, Iss. 30; NIHR Journals Library; Perth Scotland; 146 pages.

Huffington Post; "Jenny Olenick Dies During Wisdom Teeth Surgery, Parents Sue 'Negligent' Dentists"; *Huffington Post*; posted Dec. 15, 2011, updated Dec. 20, 2011; USA; 2 pages.

International Search Report and Written Opinion for Int'l Application No. PCT/US2018/053695, dated Nov. 30, 2018. 6 pages.

Jasinevicius et al.; "Prophylactic third molar extractions: US dental school departments' recommendations from 1998/99-2004/05"; *Quintessence International*; Feb. 2008; pp. 165-176; vol. 39, No. 2; Quintessenz Verlags-GmbH; USA; 12 pages.

Jiang et al.; "Ten-Year Insurance Claims Data Study Reveals Increase In Utilization of Surgical Extraction Codes"; *Dentail claim review*; 2006; pp. 1-2; USA; 2 pages.

Johnson et al.; "Frequency of Odontogenic Cysts and Tumors: a systematic review"; *Journal of Investigative and Clinical Dentistry*; 2014; pp. 5, 9-14; © 2013 Wiley Publishing Asia Pty. Ltd.; 6 pages.

Jorgenson et al.; "Comparison of the efficacy of a standard inferior alveolar nerve block versus articaine infiltration for invasive dental treatment in permanent mandibular molars in children: a pilot study"; *European Archives of Paediatric Dentistry*; 2020; pp. 171-177; vol. 21; published online Dec. 12, 2019 at: <https://dot.org/10.1007/s40368-019-00496-8>; © European Academy of Paediatric Dentistry; Springer Nature; 7 pages.

Jung et al.; "Radiographic evaluation of third molar development in 6-24 year olds"; *Imaging Science in Dentistry*; 2014; vol. 44; pp. 185-191; Korean Academy of Oral and Maxillofacial Radiology; Korea; 7 pages.

Kandasamy et al.; "The wisdom behind third molar extractions"; *Australian Dental Journal—Review*; 2009; pp. 284-292; vol. 54; Australian Dental Association; Australia; 9 pages.

Kim et al.; "Clinical and pathologic features related to the impacted third molars in patients of different ages: A retrospective study in the Korean population"; *Journal of Dental Sciences*; 2017; pp. 354-359; © Association for Dental Sciences of the Republic of China, China; publishing services by Elsevier B.V.; 6 pages.

Kinard et al.; "Most Patients with Asymptomatic, Disease-Free Third Molars Elect Extraction Over Retention as Their Preferred Treatment"; *Journal of Oral and Maxillofacial Surgeons*; 2010; pp. 2935-2942; vol. 68; American Association of Oral and Maxillofacial Surgeons; USA; 8 pages.

Knutsson et al.; "Dentists' decisions on prophylactic removal of mandibular third molars: a 10-year follow-up study"; *Community Dentistry and Oral Epidemiology*; 2001; pp. 308-314; vol. 29; © Munksgaard; Denmark; 7 pages.

Kolokythas, Antonia; Tongue 'Feels Different' After an Injury: What Next? *J. Oral Maxillofac. Surg.*; 2021; pp. 728-730; vol. 29; AAOMS; USA; 3 pages.

Koumaras, George M.; "What Costs are Associated With the Management of Third Molars?"; *Journal of Oral and Maxillofacial Surgeons*; Sep. 2012; pp. 8-10; vol. 70, Suppl. 1; AAOMS; USA; 3 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Kunkel et al.; "Severe third molar complications including death—lessons from 100 cases requiring hospitalization"; *Journal of Oral and Maxillofacial Surgeons*; 2007; abstract of article on pp. 1700-1706; vol. 65, No. 9; © AAOMS; © Elsevier; 1 page.
- Kunkel et al.; "Third molar complications requiring hospitalization"; *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology Endodontics (OOOOE)*; Sep. 2006; pp. 300-306; vol. 102; ; Elsevier Inc.; 7 pages.
- Lacerda-Santos, Jhonatan Thiago, et al. "Prevalence of second molar external root resorption caused by mandibular third molars: A CBCT study." *Gen. Dent* 71 (2023): 58-63. (1 page abstract).
- Lakhani et al.; "Anterior arch crowding—a possible predictor for mandibular third molar impaction"; *J. Pub. Med. Coll. Abbottabad*; Jan.-Mar. 2011; abstract of article on pp. 63-65; vol. 23, No. 1; MedGen; USA; 1 page.
- Larson, et al.; "The Effect of Removing the True Dental Follicle on Premolar Eruption in the Dog"; *Archives of Oral Biology*; 1994; pp. 271-275; vol. 39, No. 4; © 1994 Elsevier Science Ltd.; Great Britain; 5 pages.
- Laskin, Daniel M.; "Indications and contraindications for removal of impacted third molar"; *Dental Clinics of North America*; Oct. 1969; pp. 919-928; vol. 13, No. 4; USA; 11 pages.
- Lauesen et al.; "Association between third mandibular molar impaction and degree of root development in adolescents"; *Angle Orthodontist*; 2013; pp. 3-9; vol. 83, No. 1; The EH Angle Education and Research Foundation, Inc.; 7 pages.
- Levin, David; "Nipping Wisdom Teeth in the Bud"; *Magazine of the Tufts University Dental Alumni Associations*; Fall 2013; pp. 1-4; Tufts School of Dental Medicine; USA; 4 pages.
- Li et al.; "External root resorption in maxillary and mandibular second molars associated with impacted third molars: a cone-beam computed tomographic study"; *Clinical Oral Investigations*; 2019; pp. 4195-4203; vol. 23; Springer; 9 pages.
- Libdy et al.; "The ability of orthodontists and maxillofacial surgeons in predicting spontaneous eruption of mandibular third molar using panoramic serial radiographs"; *Dental Press Journal of Orthodontics*; Jul.-Aug. 2020; pp. 68-74; vol. 25, No. 4; © 2020 Dental Press Journal of Orthodontics; 7 pages.
- Ling et al.; "Which procedure is better—germectomy or surgical removal of mandibular third molars"; *International Journal of Oral & Maxillofacial Surgery*; abstract; Mar. 2017; 110-111; vol. 46, Suppl. 1; Elsevier Inc.; 2 pages.
- Liverpool Reviews and Implementation Group (LRiG); "Prophylactic removal of impacted third molars"; *Final Protocol*; Apr. 1, 2016; pp. 1-23; Liverpool Reviews and Implementation Group (LRiG) University of Liverpool; England; 23 pages.
- Lloro et al.; "The Incidence of Dental Needs During Isolated Missions Compared to Non-isolated Missions: A Systematic Review and Implications for Future Prevention Strategies"; *Military Medicine*; Mar./Apr. 2019; pp. e148-e155; vol. 184; Association of Military Surgeons of the United States; 8 pages.
- Luyen et al.; "Microwave Ablation at 10.0 GHz Achieves Comparable Ablation Zones to 1.9 GHz in Ex Vivo Bovine Liver"; *IEEE Transactions on Biomedical Engineering*; Jun. 2014; pp. 1702-1710; vol. 61, No. 6; IEEE; 9 pages.
- Ma, Yuecui, Daogui Mu, and Xiangxin Li. "Risk factors for root resorption of second molars with impacted third molars: A meta-analysis of CBCT studies." *Acta Odontologica Scandinavica* 81.1 (2023): 18-28. (1 page abstract).
- Magraw et al.; "Prevalence of Visible Third Molars in the United States Population: How Many Individuals Have Third Molars?"; *J. Oral Maxillofac. Surg.* 2016; pp. 13-17; vol. 74, No. 13-17; American Association of Oral and Maxillofacial Surgeons; USA; 5 pages.
- Marchiori et al.; "Initial third molar development is delayed in jaws with short distal space: An early impaction sign?"; *Archives of Oral Biology*; 2019; pp. 1-8; vol. 106; Elsevier Ltd.; 8 pages.
- Marchiori et al.; "Third-molar mineralization as a function of available retromolar space"; *Acta Odontologica Scandinavica*; 2016; pp. 509-517; vol. 74, No. 7; Taylor and Francis Group; 10 pages.
- Marciani, Robert D.; "Third Molar Removal: An Overview of Indications, Imaging, Evaluation, and Assessment of Risk"; *Oral and Maxillofacial Surgery Clinic of North America*; 2007; pp. 1-13; vol. 19; Elsevier Saunders; USA; 13 pages.
- Markiewicz et al.; "Corticosteroids Reduce Postoperative Morbidity After Third Molar Surgery: A Systemic Review and Meta-Analysis"; *PlumX Metrics*; Sep. 2008; abstract of article on pp. 1881-1894; vol. 66, No. 9; Elsevier; 2 pages.
- Marks et al.; "Regional control by the dental follicle of alterations in alveolar bone metabolism during tooth eruption"; *J. Oral. Pathol.*; 1987; pp. 164-169; John Wiley & Sons Ltd.; USA; 6 pages.
- Marks, SC Jr.; "The basic and applied biology of tooth eruption"; *Connect Tissue Research*; 1995; abstract of article on p. 149-157; vol. 32, No. 1-4; Medline; 1 page.
- Masson, Gabrielle; "Number of US dentists per specialty: 2021 vs. 2022"; *Becker's Dental + DSO Review data*; 1 webpage; Feb. 8, 2022; article available via Becker's Dental + DSO webpage; 1 page.
- Matzen et al.; "Radiographic signs of pathology determining removal of an impacted mandibular third molar assessed in a panoramic image or CBCT"; *British Institute of Radiology (BIR) Publications*; Oct. 25, 2016; pp. 1-17; vol. 26, No. 1; The British Institute of Radiology; Britain; 17 pages.
- Maung et al.; "Near-IR Imaging of Thermal Changes in Enamel during Laser Ablation"; *Proc SPIE Int Soc. Opt. Eng.*; Mar. 5, 2010; pp. 1-12; vol. 7546, No. 1.; www.ncbi.nlm.nih.gov/pmc/articles/PMC3175377; 12 pages.
- Mazur et al.; "Clinical Indications to Germectomy in Pediatric Dentistry: A Systematic Review"; *International Journal of Environmental Research and Public Health*; 2022; pp. 1-13; www.mdpi.com/journal/ijerph; Basel, Switzerland; 13 pages.
- Mcardle et al.; "Distal cervical caries in the mandibular second molar: an indication for the prophylactic removal of third molar teeth? Update"; *British Journal of Oral and Maxillofacial Surgery*; 2014; pp. 185-189; vol. 52; Elsevier Ltd.; 5 pages.
- Mcardle et al.; "The effects of Nice guidelines on the management of third molar teeth"; *British Dental Journal*; 2012; pp. 1-7; vol. 213, E8; Macmillan Publishers Limited; Britain; 7 pages.
- Mehlisch et al.; "Effects of a sclerosing agent on odontogenesis"; *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology Journal*; May 1971; pp. 723-730; vol. 31, No. 5; 11 pages.
- Mercier et al.; "Risk and benefits of removal of impacted third molars"; *J. Oral Maxillofac. Surg.* 1992; pp. 17-27; vol. 21; Elsevier Inc.; 11 pages.
- Mohan, et al.; "Prescription of Panoramic Radiographs in Children Using Age-based Prevalence of Dental Anomalies and Pathologies"; *Research Article*; accessed at least as early as Jun. 21, 2023; USA; 27 pages.
- Monaco et al.; "Incidence of Delayed Infections After Lower Third Molar Extraction"; *Journal of Environmental Research and Public Health*; 2022; pp. 1-7; <https://doi.org/10.3390/ijerph19074028>; MDPI; Basel, Switzerland; 7 pages.
- National Health Service (NHS); "Overview-Wisdom tooth removal"; NHS webpage; May 17, 2021; NHS, UK; available online at: <https://www.nhs.uk/conditions/wisdom-tooth-removal/>.
- Navy, Department of the; "Bumed Instruction"; Department of the Navy; 2010; pp. 1-22; Navy Medicine Website; USA; 22 pages.
- Niedzielska et al.; "Panoramic radiographic predictors of mandibular third molar eruption"; *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology (OOOOE)*; Aug. 2006; pp. 154-158; vol. 102, No. 2; Mosby, Inc.; 5 pages.
- Nivedita et al.; "Prophylactic extraction of non-impacted third molars: is it necessary?"; *Minerva Stomatologica*; Dec. 2019; Abstract only of article in vol. 68, No. 6, pp. 297-302; *Endizioni Minerva Medica*; 1 page.
- Norton et al.; "Concordance between Clinical Practice and Published Evidence Findings from The National Dental Practice-Based Research Network"; *JADA*; Jan. 2014; pp. 1-17; vol. 145, No. 1; Elsevier Inc.; USA; 17 pages.
- Nunn et al.; "Retained Asymptomatic Third Molars and Risk for Second Molar Pathology"; *Journal of Dental Research*; 2013; pp. 1095-1099; vol. 92, No. 12; International & American Associations for Dental Research; USA; 5 pages.

(56)

References Cited

OTHER PUBLICATIONS

O'Neil, Raymund; "The impacted mandibular third molar, indications and assessment for surgery"; *Apex*; Autumn 1968; pp. 8-11; vol. 2, No. 9; USA; 6 pages.

Oenning et al.; "Mesial Inclination of Impacted Third Molars and Its Propensity to Stimulate External Root Resorption in Second Molars—A Cone-Beam Computed Tomographic Evaluation"; *J. Oral Maxillofac. Surg.*; 2015; pp. 379-386; vol. 73; American Association of Oral and Maxillofacial Surgeons; USA; 8 pages.

Offenbacher et al.; "What Are the Local and Systemic Implications of Third Molar Retention"; *J. Oral Maxillofac. Surg.*; 2012; pp. S58-S65; vol. 70, Suppl. 1; American Association of Oral and Maxillofacial Surgeons; USA; 8 pages.

Office Action (Notice of Allowance and Fees Due (PTOL-85)) dated Oct. 11, 2023 for U.S. Appl. No. 17/515,360 (pp. 1-9).

Okunev, et al.; "Trends in national opioid prescribing for dental procedures among patients enrolled in Medicaid"; *JADA*; Aug. 2021; pp. 622-631; vol. 152, No. 8; <http://jada.ada.org>; ADA; USA; 10 pages.

Orenuga et al.; "A radiographic study of third molar crown development in a group of Nigerian children"; *Department of Child Dental Health; Pediatric Dental Journal*; 2011; pp. 107-115; vol. 21, No. 2; Elsevier Ltd.; 9 pages.

Orhan et al.; "Radiographic evaluation of third molar development in relation to chronological age among Turkish children and youth"; *Forensic Science International*; 2007; pp. 46-51; vol. 165; Elsevier Ireland Ltd.; 6 pages.

Osensa; "Fiber Optic Temperature Sensors;" Product description; Osensa website; at least as early as May 30, 2020; published online at: www.osensa.com; Burnaby, BC, Canada; 3 pages.

Palmer, Craig; "CMS, ADA reports measure post-recession dental economy"; *ADA News*; Jan. 21, 2013; vol. 44 No. 2; pp. 1 and 9; USA; 2 pages.

Palmer, Craig; "Dental spending growth projected through 2021"; *ADA News*; Jun. 18, 2012; p. 8; ADA; USA; 1 page.

Palmer, Craig; "Dental spending growth resumes"; *ADA News*; Jan. 16, 2012; pp. 1 and 8; vol. 43, No. 2; Jan. 16, USA; 2 pages.

Palmer, Craig; "Growth in dental spending expected to slow in 2009"; *ADA News*; Feb. 26, 2009; www.ada.org/news/1103.aspx; USA; 1 page.

Passarelli, et al.; "Quality of Life of Patients with Mandibular Third Molars and Mild Pericoronitis. A Comparison between Two Different Treatments: Extraction or Periodontal Approach"; *Antibiotics*; Apr. 29, 2020; *MDPI*; vol. 9; pp. 1-5; doi: 10.3390/antibiotics9050222; *MDPI*; 5 pages.

Patent Cooperation Treaty; "International Preliminary Report on Patentability" for International application No. PCT/US2020/036508; May 17, 2021; 6 pages.

Patent Cooperation Treaty; "International Preliminary Report on Patentability" for International application No. PCT/US2020/036705; Dec. 7, 2021; 10 pages.

Patent Cooperation Treaty; "International Preliminary Report on Patentability" for International application No. PCT/US2020/057383; Apr. 3, 2023; 7 pages.

Pearson, Anthony A.; "The early innervation of the developing deciduous teeth"; *Journal of Anatomy (J. Anat.)*; 1977; pp. 563-577; vol. 123, No. 3; *J. Anat.*; 15 pages.

Petsos, et al.; "Five-Years Periodontal Outcomes of Early Removal of Unerupted Third Molars Referred for Orthodontic Purposes"; *Journal of Oral and Maxillofacial Surgery*; 2021; pp. 520-531; vol. 79; © 2020 AAOMS; USA; 12 pages.

Philips et al.; "How Predictable Is the Position of Third Molars Over Time?"; *Journal of Oral and Maxillofacial Surgery*; 2012; pp. S11-S14; vol. 70, Suppl. 1; American Association of Oral and Maxillofacial Surgeons; USA; 4 pages.

Phillips et al.; "Recovery after third-molar surgery: The effects of age and sex"; *American Journal of Orthodontics and Dentofacial Orthopedics*; Dec. 2010; pp. 700.e1-700.e8; vol. 138, No. 6; American Association of Orthodontists; USA; 8 pages.

Piecuch, Joseph F.; "What Strategies Are Helpful in the Operative Management of Third Molars?" *Journal of Oral and Maxillofacial Surgery*; 2012; pp. 25-32; vol. 70, No. 9, Suppl. 1; American Association of Oral and Maxillofacial Surgeons; USA; 8 pages.

Pogrel, M. Anthony; "What Are the Risks of Operative Intervention?"; *Journal of Oral and Maxillofacial Surgery*; 2012; pp. S33-S36; vol. 70, Suppl. 1; American Association of Oral and Maxillofacial Surgeons; USA; 4 pages.

Portalatin Ariana; "Employment levels for 6 dental roles"; *Becker's Dental + DSO Review; Staffing Issues*; Jun. 29, 2023; Becker's Dental website, 1 page.

Poswillo, David; "Surgical options for third molars: a review"; *Journal of the Royal Society of Medicine*; Dec. 1981; pp. 911-913; The Royal Society of Medicine; England; 3 pages.

Prakash et al.; "Introduction to microwave tumour ablation special issue"; *International Journal of Hyperthermia*; 2016; pp. 1-3; vol. 33, No. 1; Taylor & Francis Group; 3 pages.

Prakash et al.; "An Optimal Sliding Choke Antenna for Hepatic Microwave Ablation"; *IEEE Transactions on Biomedical Engineering*; Oct. 2009; pp. 2470-2476; vol. 56, No. 10; IEEE; 7 pages.

Rafetto, Louis K.; "Managing Impacted Third Molars"; *Oral Maxillofacial Surgery Clinics of North America*; Aug. 2015; pp. 363-371; vol. 27, No. 3; Cross Mark (Elsevier); USA; 9 pages.

Rasmusson et al.; "Bisphosphonate Associated Osteonecrosis of the Jaw: An Update on Pathophysiology, Risk Factors, and Treatment"; *International Journal of Dentistry*; 2014; pp. 1-10; Hindawi Publishing Corporation; 10 pages.

Renton et al.; "What Has Been the United Kingdom's Experience With Retention of Third Molars?"; *Journal of Oral and Maxillofacial Surgery*; 2012; pp. S48-S57; vol. 70; American Association of Oral and Maxillofacial Surgeons; USA; 10 pages.

Retrouvey et al.; "Dental Development and Maturation, from the Dental Crypt to the Final Occlusion"; *Pediatric Bone* second edition; 2012; pp. 83-108; Elsevier Inc.; 26 pages.

Richardson, Margaret E.; "The Etiology and Prediction of Mandibular Third Molar Impaction"; *The Angle Orthodontist*; Jul. 1, 1977; pp. 165-172; vol. 47, No. 3; Allen Press; 8 pages.

Ricketts, Robert Murray; "Studies leading to the practice of abortion of lower third molars"; *Dental Clinics of North America*; Jul. 1979; pp. 393-411; vol. 23, No. 3; ScienceDirect; USA; 20 pages.

Ruvo et al.; "The Impact of Delayed Clinical Healing After Third Molar Surgery on Health-Related Quality-of-Life Outcomes"; *Journal of Oral and Maxillofacial Surgery*; 2005; pp. 929-935; vol. 63; American Association of Oral and Maxillofacial Surgeons; USA; 7 pages.

Ryalat et al.; "Impaction of lower third molars and their association with age: radiological perspectives"; *BMC Oral Health*; 2018; pp. 1-5; vol. 18, No. 58; CrossMark; USA; 5 pages.

Sakhdari, Shirin, et al. "Frequency and Severity of Second Molar External Root Resorption Due to the Adjacent Third Molar and Related Factors: A Cone-Beam Computed Tomography Study." *Frontiers in Dentistry* 18 (2021), 7 pages.

Salagnac, J.M.; "Is the recommendation for the enucleation or the extraction of 3rd molars in subjects during or at the end of dento-facial orthopedic treatment always justified? The viewpoint of a practitioner after 40 years of orthodontic practice"; *Journal Dentofacial Anom Ortho*; 2014; pp. 1-14; vol. 17, No. 403; *JDAO Journal*; Great Britain; 14 pages.

Saltzman, Wendy; "Three Georgia boys die unexpectedly after wisdom tooth extractions"; *CBS Atlanta News* article; Apr. 26, 2012; *CBS Atlanta News*; USA; 3 pages.

Sanchez-Labrador et al.; "Autogenous Dentin Graft in Bone Defects after Lower Third Molar Extraction: A Split-Mouth Clinical Trial"; *Materials*; Jul. 10, 2020; pp. 1-17; doi: 10.3390/ma13143090; www.mdpi/journal/materials; *MDPI*; 17 pages.

Santosh, P.; "Impacted Mandibular Third Molars: Review of Literature and a Proposal of a Combined Clinical and Radiological Classification"; *Annals of Medical and Health Sciences Research*; Jul.-Aug. 2015; pp. 1-6; vol. 5, No. 4; *Annals of Medical and Health Sciences Research*; USA; 6 pages.

Sarica et al.; "A Retrospective Study: Do All Impacted Teeth Cause Pathology?"; *Nigerian Journal of Clinical Practice*; 2019; 527-533; vol. 22; Wolters Kluwer—Medknow; 7 pages.

(56)

References Cited

OTHER PUBLICATIONS

Sarikov et al.; "Inferior Alveolar Nerve Injury after Mandibular Third Molar Extraction: A Literature Review"; *Journal of Oral and Maxillofacial Research*; 2014; pp. 1-15; vol. 5, No. 4; *Journal of Oral and Maxillofacial Research*; 15 pages.

Schroeder et al.; "Estimated Cumulative Incidence of Wisdom Teeth Extractions in Privately Insured US Patients"; *Frontiers in Dental Medicine—brief research report*; Jul. 8, 2022; pp. 1-4; vol. 3, Article 937165; www.frontiersin.org; 4 pages.

Schroeder et al.; "Association of Opioid Prescriptions From Dental Clinicians for US Adolescents and Young Adults with Subsequent Opioid Use and Abuse"; *JAMA Internal Medicine*; 2018; pp. E1-E8; American Medical Association; USA; 8 pages.

Selinger et al.; "Inhibition of tooth development with a sclerosing agent, sodium tetradecyl sulfate"; *Journal of Dental Research*; 1966; pp. 236-242; vol. 45, No. 2; Sage Publications USA; 8 pages.

Shilling, Erik; "Patient gets \$9.8 million in wisdom tooth suit"; *USA Today*; Oct. 12, 2012; USA; 1 page.

Shin et al.; "Prevalence of pathologies related to impacted mandibular third molars"; *Springer Plus*; published online Jun. 29, 2016; pp. 1-5; vol. 5, No. 915; Springer Open; 5 pages.

Extended European Search Report issued in App. No. EP20960099, dated Jan. 23, 2024, 7 pages.

Office Action (Non-Final Rejection) dated Jan. 8, 2024 for U.S. Appl. No. 18/250,534 (pp. 1-14).

* cited by examiner




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5





FIG. 6






FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27




FIG. 28




FIG. 29

THERAPEUTIC TOOTH BUD ABLATION

The present application is a continuation of U.S. patent application Ser. No. 15/886,729, filed Feb. 1, 2018. U.S. patent application Ser. No. 15/886,729 is a continuation of U.S. patent application Ser. No. 15/694,794, filed Sep. 2, 2017. U.S. patent application Ser. No. 15/694,794 is a continuation of U.S. patent application Ser. No. 13/093,842, filed Apr. 26, 2011 (which issued as U.S. Pat. No. 9,827,068 on Nov. 28, 2017). U.S. patent application Ser. No. 13/093,842 is a continuation of U.S. patent application Ser. No. 12/863,183, filed Jul. 15, 2010 (which issued as U.S. Pat. No. 9,402,693 on Aug. 2, 2016). U.S. patent application Ser. No. 12/863,183 is a national stage filing under 35 U.S.C. 371 of International Application No. PCT/US10/34259, filed May 10, 2010. International Application No. PCT/US10/34259 is an international application claiming the benefit under 35 U.S.C. Section 119(e) of U.S. Provisional Patent Application Ser. No. 61/177,143, filed May 11, 2009. The present application is also a continuation of U.S. patent application Ser. No. 15/694,794, filed Sep. 9, 2017. The present application is also a continuation of U.S. patent application Ser. No. 15/694,791, filed Sep. 9, 2017. The present application is also a continuation of U.S. patent application Ser. No. 13/093,844, filed Apr. 26, 2011. The present application is based on and claims priority from these applications, the disclosures of which are hereby expressly incorporated herein by reference in their entirety.

COPYRIGHT NOTICE

A portion of the disclosure of this patent document contains material that is subject to copyright protection. The copyright owner has no objection to the facsimile reproduction of the patent disclosure as it appears in the Patent and Trademark Office patent files or records, but otherwise reserves all copyright rights whatsoever.

TECHNICAL FIELD

Described herein are a tooth bud ablation (TBA) procedure and a tooth bud ablation (TBA) system.

BACKGROUND ART

Approximately 3.5% of the total \$100 billion spent on dental care in the United States in 2008 was for traditional surgical removal of third molars (i.e. “wisdom teeth” extractions), including the associated costs of imaging, sedation, and resulting complications. Traditional surgical removal of third molars, however, is a highly invasive, painful, and complication-ridden procedure. Further, third molar extraction represents the only procedure in the United States and Europe where it is considered “normal” to subject patients of any age group to such a highly invasive prophylactic surgery that carries significant lifelong risks for the excision of asymptotic or non-pathologic tissue. Dental practitioners (e.g. general dentists, pediatric dentists, and oral surgeons) have been trained to remove children’s wisdom teeth (third molars) before the wisdom teeth cause problems, but this surgery carries significant pain, risks, and costs.

The main problem associated with third molar tooth extractions—aside from the pain inflicted—is the serious risk of complications associated with such an invasive procedure. Each year “more than 11 million patient days of ‘standard discomfort or disability’—pain, swelling, bruising, and malaise—result post-operatively, and more than

11,000 people suffer permanent paraesthesia—numbness of the lip, tongue, and cheek—as a consequence of nerve injury during the surgery. At least two thirds of these extractions, associated costs, and injuries are unnecessary, constituting a silent epidemic of iatrogenic injury that afflicts tens of thousands of people with lifelong discomfort and disability.”

If you interview people under the age of 40 and ask them what has been the most invasive surgical procedure they have personally experienced (that is not trauma related), there is a greater than 90% chance that it will be their “wisdom teeth” extraction. The current standard of care in America for “managing” third molars (e.g. “wisdom teeth”) in adolescents and young adults is generally to have all four third molars extracted once they are formed, unless it is absolutely clear that these teeth will erupt normally. General dentists and oral surgeons alike are complicit in their belief that third molars generally should be extracted because not all will erupt normally, thus causing future pathology.

Each year, an estimated 10 million third molar tooth extractions account for over 92% of all teeth extracted for patients under the age of 40. This represents surgery on approximately 5 million people each year at an estimated cost of over \$2.5 billion for third molar extraction fees alone in the United States. When IV sedation fees, X-ray imaging expenses, post-op medications, and unplanned post-operative expenses associated with treating complications are added in, the true United States health care cost is estimated to be well over \$3.5 billion. In addition to fee inflation, it has been shown that “upcoding” of wisdom teeth extraction (i.e. using an insurance code for payment of a higher fee than is clinically justified) has become an increasing problem for insurers. Insurance claims patterns clearly show that this procedure is largely treated as an elective procedure. The average annual income per oral surgeon has been estimated to be approximately \$500,000 for third molar extraction fees alone. Insurance companies have historically reported that reimbursement for third molar extractions has been the highest reimbursed surgical procedure—even higher than hysterectomies in years when medical insurance used to pay for both procedures.

The market demographics and associated expenses are compelling. Over 77% of children at age 6 have all four third molar tooth buds radiographically detectable on routine panoramic X-rays (a type of volume scan). Over 90% of all teenagers in the United States have at least one third molar that will fully form. A typical cost for an oral surgeon to remove all four third molars on a teenager is generally \$2,000 to \$2,500 per patient once the teeth have at least partially formed—but before they have erupted—including the cost of IV sedation, consultations, and X-ray imaging costs.

There has been considerable controversy for the past fifty years regarding prophylactic extraction of third molars. A number of leading authorities have objectively tried to demonstrate that prophylactic extraction is a waste of health-care dollars, citing studies that indicate there is no objective scientific evidence for such a procedure, while other groups vigorously argue that prophylactic extraction in the teens and early adult years greatly eliminates more serious problems later in life and is worth the cost and risk.

An important question to ask is, “What happens if no prophylactic third molar extractions occur?” For instance, “as many as 22% of all emergency department visits” at a United States military support facility were related to dental problems, most of which were third-molar specific. In third-world countries, where prophylactic extraction of wisdom teeth is simply not performed, a high percentage of

patients will present with acute infections, decay, gum disease and other problems later in life. In Jordan—where prophylactic extraction is not performed—46% of adult patients had pathology (decay, infection, bone loss, etc.) detectable on their third molars on routine X-rays and volume scans. Numerous studies show that third molars are hard to clean, generally do not erupt fully, and are the single most-likely teeth to have problems associated with them.

Routine panoramic X-rays of adults taken during a random two-week period are shown in FIGS. 1 and 2. These X-rays show the examples of the range of problems that adult patients experience when they have third molars that are not extracted at an early age, including advanced decay and gum infections. For example, FIG. 1 shows a 48-year-old patient with both upper third molars present. There is a gum infection around both third molars that has caused 90% of the bone on the distal side of the second molars to be destroyed. In order to save the first molars, extraction of the second and third molars on the upper-arch will be necessary. FIG. 2 shows another example in which a 36-year-old patient has all four third molars present. The upper third molars are hyper-erupting because they have no opposing teeth to occlude against. They will eventually need to be extracted. The lower third molars are horizontally impacted and show no signs of infection, but if they become infected, then the patient will almost certainly lose the adjacent second molars because of the bone damage that will occur.

The problem all practitioners face is that it is practically impossible to tell in advance which impacted wisdom teeth will ultimately cause future pathology. The reality is that most wisdom teeth (well over 50%) are surgically extracted prophylactically with no real knowledge that they will actually cause future pathology.

If pathology appears in patients over the age of 40, however, the stakes are much different. According to two prospective studies in the United States, in 1997 10.5% and in 2002 17.3% of patients requiring third molar extractions were over the age of 40. If a patient is presenting later in life to have one or more third molar extracted, it is because active pathology has been diagnosed, making surgery no longer elective. The attendant complication rates are not just higher, but these patients were categorized as “very high risk patients” for surgery. These studies concluded, “[t]he risk to patients and to the profession can be dramatically reduced by considering early removal of abnormal third molars” and “based on our experience, we propose extraction of third molars during adolescence when the X-ray indicates normal eruption cannot be expected due to lack of space or an abnormal position.”

The occurrence of post-operative complications is generally considered to be over 15% by most independent researchers. For instance, the formation of long-term periodontal pockets on the distal surfaces of second molars that results in gum disease, infection, and eventual second molar tooth loss is estimated to be over 10% due to the damage and poor bone morphology that result from third molar extraction surgery. The incidence of post-operative infections and “dry sockets” is generally accepted to be over 15%. Temporary paraesthesia due to damage to the mandibular nerve or the lingual nerve is over 10%, with residual permanent numbness of the lip or tongue present in approximately 1.5% of all patients. Recently, it has been concluded that approximately 23% of all cases of long-term Temporomandibular Joint (“TMJ”) dysfunction and chronic joint pain are attributable to third molar extraction surgeries.

Malpractice claims against dental practitioners relating to third molar extractions are at an all time high. Litigation for

residual TMJ problems is increasing; in 2002 a North Carolina jury awarded \$5 million in damages to a patient with TMJ pain following third molar extractions. The incidence of litigation over permanent numbness of the lip has dramatically increased in recent years. Malpractice claims with resulting payouts have been reported to be as high as two-thirds of all claims made against dental practitioners when nerve damage is involved.

If the wisdom teeth are not extracted in adolescence, the roots will fully form, making future extraction difficult and dramatically increasing the incidence of serious complications if surgery should later be required. The damage induced by long-standing, chronic infections in adults may necessitate the extraction not only of the third molars when they become symptomatic, but also of the adjacent second molars. Additional complications include the reduced healing response of adults as compared to adolescents, and the economic hardship induced by having to miss work. Many references indicate that prophylactic extraction of third molars in teens and young adults—in spite of the possibility of lifelong complications such as nerve damage—is justified to avoid the non-elective third molar extraction in adults over the age of 30.

Complications can be severe, even requiring hospitalization when teeth have been extracted on an out-patient basis. There have even been reports of patients who died as a direct result of wisdom tooth extractions.

As an example, FIG. 3 is an X-ray showing a 9-year-old patient with four third molar tooth buds present; three of them are in very early stages of enamel formation. The lower right third molar tooth bud does not have enamel formed yet, but will shortly. This X-ray shows an example of the early stages in which the tiny third molar tooth buds begin to form, begin to develop enamel, and finally begin to develop roots. Early signs of problems are almost always clearly evident by the time a patient is a teenager.

Once the tooth starts to form, the tooth bud starts to become encased in bone and appears to be “pushed down” into the mandible and maxilla as the child’s jaw bone grows out and around the tooth bud with age. Future surgical access becomes far more invasive as the bone encases the forming third molar. Given the basic physiology involved, early intervention is the only approach that will eliminate the complications and high costs associated with extraction of fully formed third molars later in life.

The idea of prophylactic third molar tooth bud removal is not new. In 1936, Dr. Henry supported the surgical enucleation of tooth buds, and it was again supported in the mid 70s by several practitioners using somewhat invasive surgical techniques to physically access the tooth buds and mechanically cut them out. In 1979, Drs. Gordon and Laskin used cryoprobes to enucleate third molar tooth buds in dogs. However, at the *NIH Conference On Third Molars in 1979* it was concluded that “[a]lthough there are cogent reasons for early removal of third molars, the group felt that the suggested practice of enucleation of third molar tooth buds, based on predictive studies at age 7 to 9, is not currently acceptable.” (National Institutes of Health—*Removal Of Third Molars Consensus Development Conference Statement—1979*.)

Early removal of partially formed third molars (sometimes referred to as a “germectomy”) where the enamel of the crown has completely formed but less than one-third of the root length has formed, is demonstrated to be somewhat less invasive and carries no demonstrated long-term complications or risks associated with early-stage surgery. However, it is still highly invasive and generally requires IV

sedation of the teenage patient. The American Association of Oral & Maxillofacial Surgeon's White Paper On Third Molar Data references five studies involving over 1,100 germectomies with not a single case of a long-term complication (nerve injury, etc.) associated with the surgery. Further, since the germectomies were carried out on teenagers, there were no economic hardships induced by missing work. The White Paper understates the obvious conclusions associated with early intervention: "It does appear that early third molar removal may be associated with a lower incidence of morbidity and also less economic hardship from time off work for the patient." However, it can also be concluded that there is a tremendous conflict of interest because this paper was written by oral surgeons. To date there is still no measurable shift by dental practitioners to change the way in which third molars are screened, diagnosed, and extracted (i.e. early extraction), indicating that there is a need to fundamentally change the way this condition is being surgically managed.

There are a number of existing alternative technical approaches that can be considered for prophylactic enucleation of third molar tooth buds before the crown or root begins formation in children age 6 to 10. These technical approaches include ablation procedures using different types of ablation means. Exemplary ablation procedures include electrourge tissue ablation (rats), cryoablation (dogs), laser ablation (dogs), and the use of a scalpel (humans). All but the first three ablation procedures (microwave ablation, radio frequency ablation, and irreversible electroporation) have significant limitations due to being highly invasive, high in cost, requiring cumbersome equipment, or due to the limited means of mechanical access in the oral cavity. Nor do these ablation procedures offer the potential for real-time feedback control to contain collateral tissue damage. To date, the only documented trial of any form of tooth bud ablation procedure utilizing ablation technology that is currently used in mainstream medicine is cryoablation (although preliminary animal trials have been completed using electro-surgical power and lasers).

The article entitled "Selectively Preventing Development Of Third Molars In Rats Using Electro-surgical Energy" by Silvestri et al. describes a pilot study that tests the hypothesis that third molars can be selectively prevented from developing. To test the hypothesis, a study was conducted in which thirty-three neonate rats received electro-surgical energy to the mucosal surfaces of one of their maxillary tuberosities. In this study, guides (insulating plastic positioning devices that housed the electro-surgical probes) were used. The guides were fabricated using the mouths of euthanized rat pups of the same age as the rats that were to be treated as a mold for creating the guides. Then, the electro-surgical probe placed so that its stainless steel tip extended less than 1.0 mm beyond the plastic positioning device to ensure contact with the external surface of the oral mucosa of the maxillary tuberosity. Finally, when in position, the rat pups received a single, unilateral, momentary pulse of monopolar electro-surgical energy to the external surface of the gum tissue of their maxillary tuberosity regions. It should be emphasized that this surface application of electro-surgical energy acted first to unnecessarily kill the overlying gum tissue, then bore a hole through the gum tissue, and otherwise damage not only the tooth buds, but other nearby tissue. The rats were cared for, but after the experimental period, were euthanized to determine the effectiveness of the procedure. The results were that ten rats showed no intra-oral or radiographic evidence of third molar development (and most of these rats subsequently developed

palatal deformities), and six developed smaller-than-normal third molars. The conclusion was that maxillary third molars could be selectively prevented from developing in rat pups at or near the time of tooth bud initiation. It was recognized, however, that electro-surgical energy was too powerful and uncontrollable to reliably confine its damage to only the tooth-forming tissues.

SUMMARY OF THE INVENTION

Described herein is a method for creating a tooth bud ablation system including a custom surgical stent and an ablation probe tip for use in a tooth bud ablation procedure that results in the agenesis of a tooth in a patient. The method preferably includes the following steps: (a) measuring a three-dimensional location and volume of a tooth bud of the patient to obtain three-dimensional location measurements and volume measurements; (b) obtaining the ablation probe tip, the ablation probe tip having a shaft and a center of ablation, the center of ablation positioned on the shaft; (c) calculating a pre-defined angle to guide the ablation probe tip so that the center of ablation is within the tooth bud, the calculations of the pre-defined angle being based on the three-dimensional location measurements and volume measurements; (d) calculating a pre-defined depth to limit the depth of the ablation probe tip so that the center of ablation is within the tooth bud, the calculations of the pre-defined depth being based on the three-dimensional location measurements and volume measurements; and (e) creating the custom surgical stent with at least one surgical guide corresponding to at least one tooth bud surgical site, the at least one surgical guide having guiding structure, the at least one surgical guide being a passageway through the surgical stent, the passageway being sized to guide placement of the ablation probe tip only at the pre-defined angle, the at least one surgical guide having mechanical stop structure to limit the depth of the ablation probe tip to the pre-defined depth.

Described herein is a tooth bud ablation procedure that results in tooth agenesis, including the steps of: (a) physically seating a custom surgical stent having at least one surgical guide so the at least one surgical guide corresponds to at least one tooth bud surgical site; (b) using the at least one surgical guide, making a surgical access path at the at least one tooth bud surgical site; (c) using the at least one surgical guide, guiding placement of an ablation probe tip having a center of ablation so that the center of ablation is in the middle of a tooth bud at the at least one tooth bud surgical site; and (d) at least partially ablating at least one tooth bud.

Described herein is a tooth bud ablation system for use in a tooth bud ablation procedure that results in tooth agenesis, the system including: (a) a custom surgical stent with at least one surgical guide corresponding to at least one tooth bud surgical site; (b) an ablation probe tip having a center of ablation; and (c) the at least one surgical guide having structure for guiding placement of the ablation probe tip so that the center of ablation is in the middle of a tooth bud by inserting the ablation probe tip through the at least one surgical guide.

Described herein is an ablation procedure including the steps of: (a) physically seating a custom surgical stent having at least one surgical guide so the at least one surgical guide corresponds to at least one lesion or tumor surgical site; (b) using the at least one surgical guide, making a surgical access path at the at least one lesion or tumor surgical site; (c) using the at least one surgical guide, guiding placement of an ablation probe tip having a center of

ablation so that the center of ablation is in the middle of a lesion or tumor at the at least one lesion or tumor surgical site; and (d) at least partially ablating at least one lesion or tumor.

Described herein is an ablation procedure including the steps of: (a) physically seating a custom surgical stent having at least one surgical guide so the at least one surgical guide corresponds to at least one lesion or tumor surgical site; (b) using the at least one surgical guide, guiding placement of an ablation probe tip having a center of ablation so that the center of ablation is in the middle of a lesion or tumor at the at least one lesion or tumor surgical site; and (c) at least partially ablating at least one lesion or tumor.

Described herein is a method for volume scanning both hard tissues and soft tissues of a patient, the method including the steps of: (a) using an impression of a material visible in a volume scan; (b) generating a volume scan in which hard tissue is visible and the impression is visible, and soft tissue being “visible” as the space between the visible hard tissue and the visible impression; and (c) providing results of the step of generating a volume scan for the purpose of manufacturing or fabricating a custom surgical stent having at least one surgical guide for guiding placement of an ablation probe tip.

Described herein is a method for simultaneous volume scanning of both hard tissues and soft tissues, the method including the steps of: (a) using a dental impression of a material visible in a volume scan; (b) physically seating the dental impression in a patient’s mouth; (c) volume scanning the patient’s mouth while the dental impression is seated therein; (d) the step of volume scanning generating a volume scan in which hard tissue is visible and the dental impression is visible, and soft tissue is “visible” as the space between the visible hard tissue and the visible dental impression; and (e) providing the results of the step of volume scanning for the purpose of manufacturing or fabricating a custom surgical stent having at least one surgical guide for guiding placement of an ablation probe tip.

Described herein is a method for manufacturing or fabricating a custom surgical stent, the method including the steps of: (a) using a volume scan image in which hard tissue is visible and a dental impression is visible, and soft tissue is “visible” as the space between the visible hard tissue and the visible dental impression; and (b) manufacturing or fabricating a custom surgical stent with at least one ablation probe tip guide for guiding at least one ablation probe tip to a pre-defined angle and depth of ablation based on information obtained from the volume scan image.

Described herein is a tooth bud ablation procedure that results in tooth agenesis, including the steps of: (a) pre-operatively taking measurements to determine a three-dimensional location of the middle of a tooth bud; (b) placing an ablation probe tip having a center of ablation so that the center of ablation is in the three-dimensional location of the middle of a tooth bud; and (c) at least partially ablating at least one tooth bud.

Described herein is a custom surgical stent for use in a tooth bud ablation procedure that results in tooth agenesis, the custom surgical stent for use with an ablation probe tip having a center of ablation, the stent including: (a) a custom surgical stent with at least one surgical guide corresponding to at least one tooth bud surgical site; (b) the at least one surgical guide having guiding structure to guide placement of an ablation probe tip at a pre-defined angle so that a center of ablation of the ablation probe tip is in the middle of a tooth bud; and (c) the at least one surgical guide having

mechanical stop structure to limit the depth of the ablation probe tip to a pre-defined depth.

The foregoing and other objectives, features, and advantages of the invention will be more readily understood upon consideration of the following detailed description of the invention, taken in conjunction with the accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying drawings are incorporated in and constitute a part of this specification.

FIG. 1 is an X-ray showing a 48-year-old patient with both upper third molars present, the X-ray being presented to show examples of the range of problems that adult patients experience when they have third molars that are not extracted at an early age.

FIG. 2 is an X-ray showing a 36-year-old patient with all four third molars present, the X-ray being presented to show examples of the range of problems that adult patients experience when they have third molars that are not extracted at an early age.

FIG. 3 is an X-ray showing a 9-year-old patient with four third molar tooth buds present; three of them are in very early stages of enamel formation, but the lower right third molar tooth bud does not yet have enamel formed.

FIG. 4 is a flow chart showing steps in preferred TBA procedures including: (1) routine screening and diagnosis; (2) pre-surgical impressions and scanning; (3) assembling a TBA surgical kit; (4) operator delivery of the TBA procedure; and (5) follow-up.

FIG. 5 is a simplified block diagram of a TBA probe system, a custom surgical stent, and a tooth bud.

FIG. 6 is a cross-sectional side view of an ablation probe tip in the process of being inserted through a surgical guide of a stent.

FIG. 7 is a cross-sectional side view of an ablation probe tip inserted through a surgical guide of a stent into the tooth bud.

FIG. 8 is a cross-sectional side view of an ablation probe tip having a linear array of temperature sensors inserted in the tooth bud.

FIG. 9 is a cross-sectional side view of an ablation probe tip ablating the tooth bud.

FIG. 10 is a cross-sectional side view of an ablation probe tip being removed from the ablated tooth bud.

FIG. 11 is a flow chart showing the steps of a TBA procedure that result in tooth agenesis.

FIG. 12 is a flowchart showing the steps that a software program for manufacturing or fabricating custom surgical stents and defining (and/or computing or calculating) the pre-determined parameter settings and/or treatment time settings.

FIG. 13 is a panoramic X-ray showing a patient whose third molar tooth buds in the #17 and #32 positions are treatable by TBA.

FIG. 14 is a pre-operative cone beam computed tomography (“CBCT”) scan of a different patient.

FIG. 15 is a series of X-rays showing successive 1.0 mm slices through both #17 and #32 in 1.0 mm increments.

FIG. 16 is a perspective view from a front corner showing a pre-operative upper-arch impression being taken of a simulated patient.

FIG. 17 is a cross-sectional view of an upper-arch impression being taken of a simulated patient.

FIG. 18 is a perspective view from above of the completed upper-arch impression.

FIG. 19 is a perspective view from above of the completed upper-arch impression, along with a stone model that will serve as a “positive” for manufacturing or fabricating of a custom surgical stent for that patient’s upper-arch.

FIG. 20 is a CBCT scan with notations showing the measurement of the angle of entry into the tooth bud.

FIG. 21 is a series of X-rays with notations showing the measurement of the lateral angle of entry.

FIG. 22 is a CBCT scan with highlights showing the computed volume of each tooth bud.

FIG. 23 is a perspective view from above of a surgical stent with two surgical guides, the stent having been manufactured or fabricated using the CBCT positioning information.

FIG. 24 is a perspective view showing topical anesthetic being applied to the base of the surgical guide.

FIG. 25 is perspective view from a front corner of a surgical stent being seated on the upper-arch of the simulated patient.

FIG. 26 is a perspective view from a front corner of a local anesthetic being injected into a tooth bud site.

FIG. 27 is a perspective view from a front corner of a tissue trocar being used to punch to the base of a tooth bud.

FIG. 28 is a perspective view from a front corner of an ablation probe tip with a mechanical (physical) stop being positioned through the surgical guide into the tooth bud.

FIG. 29 is a perspective view from a front corner of the ablation probe tip being positioned in each tooth bud through the surgical guide so that the ablation probe tip’s effective center of ablation is in the middle of each tooth bud.

DETAILED DESCRIPTION OF THE INVENTION

The highly invasive surgical procedure of extracting third molars can be completely eliminated by prophylactically eliminating the small tooth buds that will eventually form the wisdom teeth. Children age 6 to 12 will generally have radiographically detectable tooth buds with no signs of tooth formation inside the tooth bud. Third molar tooth bud agenesis (the lack of third molar formation) can only be conclusively determined by age 14. Third molar tooth buds are lying just 2.0 mm to 3.0 mm beneath the surface of the attached gingival (gum) tissue, making them accessible for rapid anesthesia and minimally invasive ablation with the correct selection of soft tissue ablation and supporting scanning and stent-manufacturing technologies.

By successfully improving existing medical technology, the highly invasive, painful, and complication-ridden procedure of traditional surgical removal of third molars (i.e. “wisdom teeth” extractions) can be replaced with a minimally invasive tooth bud ablation (TBA) procedure 70 such as that shown in FIG. 4 that is risk free, painless post-operatively, and less expensive when compared to surgical extractions.

The TBA procedure 70 (FIG. 4) and TBA system 100 (FIG. 5) for use in the TBA procedure 70 seek to achieve: (1) a minimally invasive procedure consisting of a surgical access path at a surgical site (e.g. at each tooth bud surgical site), (2) that can predictably ablate all four third molar tooth buds 120 in thirty (30) minutes or less (including time to administer anesthesia) using either microwave (“MW”) or radio frequency (“RF”) ablation, (3) that can be administered by dental practitioners under normal office conditions, (4) with direct procedure costs reduced by 25% or more, and (5) with zero risks or complications when compared to

traditional surgical extraction of fully developed third molars. It should be noted that the TBA procedure 70 is shown and described as a prophylactic third molar tooth bud ablation (TMTBA), but it is not limited thereto. For example, there may be supernumerary teeth that should not be in a patient’s mouth (e.g. there may be two teeth #5), the removal of which would not be prophylactic in nature.

One preferred advantage of the surgical phase 90 described herein is that it is a minimally invasive surgical procedure. With a minimally invasive surgical procedure design coupled with electronic feedback controls using MW and RF ablation technology to limit soft tissue damage, performing this procedure on children aged 6-12 years old takes approximately thirty (30) (or fewer) minutes, including the time to administer local anesthetics.

Another preferred advantage of the surgical phase 90 described herein is that it will not accidentally disrupt adjacent second molar tooth development, even though the formation of second molars are well under way because these tooth buds 120 have started to form before birth. The use of relatively new scanning technologies (e.g. computed tomography volume scanning such as cone beam computed tomography (CBCT) scanning and MRI volume scanning) and accurate custom surgical stents 110 to guide ablation probe tip 108 placement will eliminate the risk of accidentally disrupting the second molars by minimizing collateral tissue damage.

Summarily, the TBA procedure 70 (FIG. 4) preferably includes a screening phase 72, a pre-surgical phase 80 (also referred to as TBA pre-surgical phase 80) that includes pre-surgical scanning 82 and the assembling of a TBA surgical kit 88 (that includes pre-determined settings 105 as well as a surgical stent 110), a surgical phase 90 (also referred to as TBA surgical phase 90), and a follow-up phase 98.

A TBA system 100 (FIG. 5) is preferably used during the surgical phase 90 (shown graphically in FIGS. 6-10 and as a flow chart in FIG. 11) of the TBA procedure 70. Summarily, the TBA system 100 includes a TBA probe system 101 (including a generator 104 capable of emitting one or more types of ablation means 104', a hand piece 106, and an ablation probe tip 108) and at least one surgical stent 110 (which was manufactured or fabricated during the pre-surgical phase 80). Each stent 110 has at least one surgical guide 112 to guide the placement of the ablation probe tip 108 so that its center of ablation 130a is placed into the middle of the tooth bud 130b. This is accomplished by positioning ablation probe tip 108 through the surgical guide 112 at a pre-defined angle and depth using a mechanical relationship of the ablation probe tip 108 and the surgical guide 112 to form a “stop” therebetween. FIGS. 6-10 show (and FIG. 11 describes) the procedure of inserting the ablation probe tip 108 through the surgical guide 112 of a stent 110, ablating the tooth bud 120, and removing the ablation probe tip 108 from the ablated tooth bud 120'.

The TBA System 100

The TBA system 100 described herein is the system that is used during the surgical phase 90 of the TBA procedure 70. Some of the components (e.g. the custom surgical stent 110 and the pre-determined settings 105) used in the TBA system 100 are part of the TBA surgical kit assembled during the pre-surgical phase 80.

The TBA system 100, as shown in FIG. 5, includes a TBA probe system 101 (including a generator 104, a hand piece 106, and an ablation probe tip 108) and at least one surgical stent 110 (each stent 110 has at least one surgical guide 112 to guide (direct) the placement of the ablation probe tip 108

11

to the middle of the tooth bud **130b**). The generator **104** and the hand piece **106** may be jointly referred to as the ablation probe unit **102** (or the programmable ablation probe unit **102**). The generator **104** and hand piece **106** may be integral or functionally connected together. The generator **104** (and/or the ablation probe unit **102**) may be programmed with pre-determined parameter settings **105a** and/or treatment time settings **105b** (referred to jointly as pre-determined settings **105**). The generator **104** (and/or the ablation probe unit **102**) provides an ablation means **104'** for ablating the tooth bud **120** based on the pre-determined settings **105**. Central to the TBA system **100**, is the interaction between the ablation probe tip **108** and the surgical stents **110** (and specifically the surgical guides **112**).

Generator **104**

The generator **104** provides the ablation means **104'** suitable for ablating a tooth bud **120** during the surgical phase **90** of the TBA procedure **70**. MW energy and RF energy are discussed as exemplary preferred ablation means **104'**. Another alternative preferred ablation means **104'** is irreversible electroporation because it has subsecond activation times that can reduce collateral tissue damage. Yet another alternative preferred ablation means **104'** include, but are not limited to, cryoablation, ultra-high intensity ultrasound, laser, chemical, thermal or hot tip (e.g. a tip having any source of heat including, but not limited to, a light bulb, a soldering iron, or steam heat), and/or mechanical means. These ablation means **104'** may also be combined either simultaneously or consecutively. It should also be noted that other known and yet-to-be-developed ablation means **104'** may also be used. It should be noted that although discussed primarily in terms of MW and RF, unless specifically set forth otherwise, the use of other ablation means **104'** is possible.

The generator **104** (alone or as part of an ablation probe unit **102**) may be programmed by the operator and/or at the laboratory and/or factory and may be accomplished automatically or manually. The programming of the generator **104** may include programming at least one pre-determined setting **105**.

The following bulleted points are exemplary details and/or features that may be incorporated in preferred generators **104**.

Preferred generators **104** may be multi-use devices designed as 110V counter-top units.

Preferred generators **104** may be MW/RF generators with output levels determined initially through finite element analysis models or experimentally derived functions that exist for tumor ablation.

Preferred generators **104** (and/or ablation probe units **102**) may have operator input mechanisms (e.g. knobs, dials, key pads, keyboards, I/O interfaces, connections to the internet, or other means for inputting or programming) in which the operator inputs (or allows input of) the pre-determined settings **105**.

Preferred generators **104** (and/or ablation probe units **102**) may have output mechanisms **103** (e.g. a display or audio) for providing setting feedback (e.g. calibration cycles and pre-determined settings **105**), warning feedback (e.g. to prevent operator mishandling), and intra-operative feedback on the progress of the procedure such as time remaining (e.g. a count down or a series of beeps to alert the operator to procedure completion) and/or temperature (e.g. to alert the operator to overheating).

Preferred output displays may be digital readout displays (that may be color and/or in a large format) that permit

12

the operator to easily see feedback intra-operatively from across a standard dental operator (approximately 6-8 feet viewing distance).

Hand Piece **106**

The hand piece **106** is the functional intermediary between the generator **104** and the ablation probe tip **108**. The hand piece **106** may be connected substantially at one end to the generator **104**. Substantially at the other end of the hand piece **106**, opposite the generator **104**, the end of the hand piece **106** (the surgical end) is adapted to accept the ablation probe tip **108**. The hand piece **106** is preferably detachable from the generator **104** (if they are not an integral unit) and the ablation probe tip **108** is preferably detachable from the hand piece **106**.

The following bulleted points are exemplary details and/or features that may be incorporated in preferred hand pieces **106**.

Preferred hand pieces **106** preferably hold or secure an ablation probe tip **108** by latching the ablation probe tip **108** into the hand piece head. In some hand pieces **106**, the ablation probe tip **108** latches into the hand piece head at an angle (e.g. a 90 degree angle). It should be noted that the terms "latching" and "latch" are used to describe any type of secure fit including, but not limited to, clipping, snapping, or holding.

Preferred hand pieces **106** preferably have a hand piece head (attached or integral) that is at an approximately 20 degree angle to the rest of the hand piece. This bend emulates a standard dental high-speed hand piece to facilitate easy access of both upper and lower surgical sites. In some preferred hand pieces **106**, the 20 degree bend can be adjusted intra-operatively to permit improved operator access to both upper- and lower-arches.

Preferred hand pieces **106** preferably are rapidly detachable from the generator **104**. Preferably the connectors are ultra-reliable for repeated removal/attachment from the generator **104**.

Preferred hand pieces **106** are preferably fully steam autoclavable. Alternative preferred hand pieces **106** are disposable or have disposable covers.

Preferred hand pieces **106** preferably have actuators to allow operator activation. The actuators may be separate from the hand pieces **106** or integral therewith. Exemplary actuators include, but are not limited to a wireless foot control or a hand-operated switch on the hand piece **106**.

The hand piece **106** may be integral with the generator **104** to form a hand-held integrated generator unit (hand-held integrated ablation probe unit).

Ablation Probe Tip **108**

One end of the ablation probe tip **108** has structure suitable for connecting it to the hand piece **106**. The ablation means **104'** flows from the generator **104** through the ablation probe tip **108** and out to a center of ablation **130a** (the focal point of the ablation). The ablation probe tip **108** is insertable through the surgical guide **112**, through the gingival tissue **122**, and into the middle of the tooth bud **130b**. The center of ablation **130a** is at the insertion end of the ablation probe tip **108** such that when the insertion end of the ablation probe tip **108** is positioned at the pre-defined angle (ϕ) and pre-defined depth (x) during the surgical phase **90**, the center of ablation **130a** substantially coincides with or overlaps the middle of the tooth bud **130b**.

The pre-defined angle (ϕ) is the angle at which the ablation probe tip's effective center of ablation **130a** is in the "middle" of the tooth bud **130b** as calculated (during the

13

pre-surgical phase **80**) as described herein or using an alternative method. The pre-defined depth (x) is the depth at which the ablation probe tip's effective center of ablation **130a** is in the "middle" of the tooth bud **130b** as calculated as described herein or using an alternative method. The phrase "middle of the tooth bud **130b**" is meant to include the three-dimensional area within the tooth bud **120** and, in particular, the three-dimensional area within the tooth bud **120** that is more towards the absolute middle point than towards the outer periphery of the tooth. The pre-defined angle (ϕ) and pre-defined depth (x) can also be referred to as the "calculated angle and depth," the "prescribed angle and depth," the "proper angle and depth," the "correct angle and depth," the "optimal angle and depth," or the "ideal angle and depth."

The ablation probe tip **108** includes a mechanical stop structure **140** (e.g. a band, protrusion, or shoulder) designed to physically limit the depth of the ablation probe tip **108** when used in conjunction with mechanical stop structure **142** (e.g. the upper surface, a protrusion on the upper surface, or a notch in the upper surface) of the surgical stent **110** and/or surgical guide **112**. In other words, the mechanical stop structure **142** of the surgical guide **112** and the mechanical stop structure **140** of the ablation probe tip **108** together limit how much of the ablation probe tip **108** can pass through the surgical guide **112** until there is a mechanical stop between the mechanical stop structure **142** of the surgical guide **112** and the mechanical stop structure **140** of the ablation probe tip **108**.

Each ablation probe tip **108** may be individually custom made (e.g. manufactured or fabricated) or may be selected from a family of ablation probe tips **108** (i.e. there may be a "family" of probe tips **108** that will cover all clinical possibilities for tooth bud diameters and depths). In the manufacturing or fabricating of the surgical stents **110**, however, the characteristics of the ablation probe tip **108** (custom made or selected) that may be taken into consideration include, for example, length, shape, angle, position of a mechanical stop structure **140**, diameter, and size, shape, and location of the center of ablation **130a**. For example, if a particular ablation probe tip **108** had mechanical stop structure **140** (shown as the bottom surface of an annular ring or shoulder in FIGS. **6-10** and **28-29**) 10.0 mm from the absolute tip of the ablation probe tip **108** (and the center of ablation **130a** is substantially adjacent to the absolute tip), but the center of ablation **130a** was only 8.0 mm from the surface of the gingival tissue **122** (shown as (x) in FIG. **6**), then the surgical guide **112** would have to be 2.0 mm thick (shown as (y) in FIG. **6**). On the other hand, if all surgical guides **112** being made by the procedure were exactly 0.5 mm thick, the ablation probe tip **108** would either have to be made or selected so that the mechanical stop structure **140** is 8.5 mm from the center of ablation **130a** of the ablation probe tip **108**. The appropriate ablation probe tip **108** preferably will result in the intra-operative placement of the effective center of ablation **130a** of the ablation probe tip **108** into the targeted middle of the tooth bud $130b \pm 0.5$ mm.

The ablation probe tips **108** may be sharp enough and/or may be strong enough so that the ablation probe tips **108** can be "self-introducing" in that the ablation probe tips **108** can be pushed through the gingival tissue **122**. Alternatively, if tissue trocars **146** (described herein) are to be used, the ablation probe tips **108** would not have to be as sharp and/or strong.

The following bulleted points are exemplary details and/or features that may be incorporated in preferred ablation probe tips **108**.

14

Preferred ablation probe tips **108** are preferably disposable (e.g. single-use).

Preferred ablation probe tips **108** may be specially designed to work with the specific ablation means **104'** produced by the generator **104**. Other preferred ablation probe tips **108** may be designed to work with multiple types of ablation means **104'** produced by the generator **104** or generators **104**.

The design of the ablation probe tip **108** may be dependent on the physics involved with transmitting ablation means **104'** through the smallest possible diameter with an ideal maximum diameter. For example, an MW/RF ablation probe tip may be designed for transmitting MW/RF energy through the smallest possible diameter with an ideal maximum diameter of 0.5 mm to 1.0 mm targeted.

The "family" of probe tips **108** may include probe tips **108** having a variety of characteristics. For example, the family might have probe tips **108** of different lengths ranging from 5.0 mm to 20.0 mm. This range would accommodate the various diameters of the tooth buds **120** and overlying gingival tissue **122** thicknesses.

Intra-operative temperature sensing (shown as being performed by a linear array of temperature sensors **144** in FIG. **8**) is preferably provided at or near the apex of the ablation probe tip **108** (assuming placement in the ideal middle of the tooth bud **130b**) and/or along the shaft **145** of the probe tip **108**. Temperature sensors **144** provide core temperatures for feedback control purposes (so that the operator can monitor the temperature and/or for software feedback control loops and emergency shutdown) and/or for safety controls to reduce or eliminate collateral tissue damage. Intra-operative tissue temperature is preferably measured, both to assure complete ablation and to prevent over-heating of tissues; this may require additional set up data or programming. If temperature sensors **144** are used, the appropriate ablation probe tip **108** preferably will result in the intra-operative placement of the effective center of ablation **130a** of the ablation probe tip **108** into the targeted middle of the tooth bud $130b \pm 1.0$ mm.

Stent **110**

The at least one custom surgical stent **110** (also referred to as a "stent **110**" or a "surgical stent **110**") has at least one surgical guide **112** (also referred to as "guides **112**" or "ablation probe tip guides **112**"). Two surgical stents **110** would be used, for example, if both upper and lower tooth buds **120** were to be ablated. The surgical stents **110** are designed to seat in a patient's mouth and may be supported by at least one tooth (a tooth-supported surgical stent), soft tissue (a soft tissue-supported surgical stent), and/or bone (a bone-supported surgical stent). If the surgical stent **110** is supported by more than one of these, it could be considered a combination-supported surgical stent. Preferred surgical stents **110** may "snap" into the mechanical undercuts inherent in the patient's erupted teeth. A surgical stent **110** would have more than one surgical guide **112** if more than one tooth bud were to be ablated on either the upper or lower jaw.

The surgical stents **110** and the guides **112** therein are used to control both the pre-defined angle (ϕ) and the pre-defined depth (x) of the ablation probe tip **108** in order to assure that the ablation probe tip's effective center of ablation **130a** is in the middle of the tooth bud $130b \pm 0.5$ mm. The pre-defined angle (ϕ) is primarily controlled by the angle of the surgical guides **112** (the passageways through the stent **110**). The pre-defined depth (x) is primarily controlled by the

interaction between the mechanical stop structure **142** of the surgical stent **110** (and/or surgical guide **112**) and the mechanical stop structure **140** of the ablation probe tip **108**. The operator inserts the ablation probe tip **108** at the entry angle (ϕ) defined by the guide **112** and to the depth (x) limited by the mechanical stop structure **140**, **142**.

The surgical guides **112** are passageways through the surgical stent (the passageways being a type of guiding structure). The pre-defined angle (ϕ) for each passageway (guide **112**) is determined by the position of the middle of the tooth bud **130b**. For example, if the middle of the tooth bud **130b** is "slightly forward" the angle (ϕ) of the passageway (guides **112**) would be "slightly forward" so that the ablation probe tip **108** is angled "slightly forward" so that the center of ablation **130a** is positioned substantially at the middle of the tooth bud **130b**. The angle (ϕ) of the passageway is determined (e.g. calculated) by the software based upon tooth bud volumes determined in pre-surgical volume scanning **82**. In addition to providing a path through which the ablation probe tip **108** accesses the gingival tissue and the tooth bud, the guides **112** may also be used to provide access for administering local anesthetic and to provide access to a tissue trocar **146** (if necessary).

In the shown preferred example, the mechanical stop structure **142** is the upper surface of the surgical stent **110** and/or surgical guide **112**. The mechanical stop structure **142** is substantially adjacent to or near the surgical guide **112**. The mechanical stop structure **142**, however, could be positioned at locations of the surgical stent **110** beyond the surgical guide **112**. Alternative preferred mechanical stop structure **142** includes a protrusion on the upper surface or a notch in the upper surface. The size and shape of the mechanical stop structure **142** is determined (calculated or designed) by a process that may be implemented as software or as a program and is based upon tooth bud volumes determined in pre-surgical volume scanning **82** as well as the length between the ablation probe tip mechanical stop structure **140** and its center of ablation **130a**. For example, if the middle of the tooth bud **130b** is 2.5 mm below the surface (determined in pre-surgical volume scanning **82**), and the available ablation probe tips **108** have a length (between their respective mechanical stop structure **140** and its center of ablation **130a**) of 2.4 mm and 2.6 mm, the process (that may be implemented by software or a program) would determine that the 2.6 mm ablation probe tip **108** is the appropriate ablation probe tip **108** (the 2.4 mm ablation probe tip **108** being too short), but that the surgical stent **110** and/or surgical guide **112** would have to be approximately 0.1 mm thick to make up the difference or the 2.6 mm ablation probe tip **108** would be able to be pushed in too far.

FIG. **12** is a flowchart showing the steps of a process (that may be implemented as one or more software program or subprograms if the shown steps are divided) that, in part, determines the pre-defined angle (ϕ) and the pre-defined depth (x) (see steps **200**, **210**, **212**, **214**, **216**, and **218**). Using this process, patient volume scans are used to accurately manufacture or fabricate custom surgical stents **110** with the correct ablation probe tip angle (ϕ) and depth (x) manufactured into them. More specifically, using this process with the volume scans will permit accurate placement of the distal surgical guides **112** onto the custom surgical stents **110** so that both angle (ϕ) of insertion and depth (x) of insertion of the ablation probe tip **108** are controlled to ± 0.5 mm, placing the ablation probe tip's effective center of ablation **130a** in the middle of the tooth bud **130b**.

The following bulleted points are exemplary details and/or features that may be incorporated in preferred stents **110**.

Preferred surgical stents **110** are preferably disposable (e.g. single-use).

Manufacturing or fabricating of the custom surgical stents **110** may be based upon Poly Vinyl Siloxane (PVS) full arch impressions of the patient's erupted teeth using either conventional lab fabrication techniques or direct-digital manufacturing or fabricating techniques. If an operator has a CBCT unit in his office, it may be possible to directly scan the PVS impressions and email the volume scan of the impression to eliminate the need to physically send them to the lab. The impression materials may include materials other than PVS and preferably will be contrast-optimized through the addition of X-ray contrast agents (such as barium or iodine) to provide optimized volume scans of the dental impression for resolving the fine surface detail of the teeth and gingival tissue **122**. This unique material would be a radiographic contrast-optimized dental impression material for high resolution X-ray CT volume scanning.

Preferred surgical stents **110** are preferably made of any appropriate material including, but not limited to, plastic, acrylic, or other nontoxic sturdy material suitable for use in a patient's mouth. One exemplary surgical stent **110** composition may be, for example, clear acrylic (polymethyl methacrylate). It should be noted that materials suitable for additive-type manufacturing (or other direct-digital manufacturing or fabricating techniques) that resulted in nontoxic sturdy stents would be preferable.

Preferred surgical stents **110** preferably have markings such as color codes or numbering clearly marking or identifying the tooth bud numbering sites.

Once the surgical stent **110** is seated onto the patient's teeth, it preferably will remain firmly in place throughout the surgical phase **90** of the TBA procedure **70**.

The operator may administer local anesthetic through the guides **112**.

Pre-Determined Settings **105**

The pre-determined settings **105** include, for example, pre-determined parameter settings **105a** and/or treatment time settings **105b** that are needed to control (provide instructions to) the generator **104** (alone or as part of an ablation probe unit **102**) to provide sufficient ablation means **104'** to ablate the tooth bud **120**, but not so much as to incur significant collateral soft tissue damage (e.g. to the gingival tissue **122**). For example, the pre-determined parameter settings **105a** might control the quantity and quality ablation means **104'** delivered to the tooth bud **120**. The actual pre-determined parameter settings **105a** will be highly dependent on the type of ablation means **104'** to be delivered. For example, MW and RF ablation means might have parameters relating to wavelength and/or frequency, hot tip ablation means might have parameters relating to temperature, chemical ablation means might have parameters relating to the strength of the chemical and how fast the chemical is flowing into the tooth bud, and mechanical ablation means might have parameters relating to speed.

The pre-determined settings **105** are determined (which includes computing, calculating, looking up, processing, or otherwise determining) by a process (that may be implemented as software or a program) based upon tooth bud volumes determined in pre-surgical volume scanning **82**. It should be noted that the pre-determined settings **105** may take into consideration factors other than tooth bud volume including, but not limited to, image recognition programs to measure tooth bud location, age and size of the patient, and

other relevant factors to successfully image the patient for the TBA procedure 70. FIG. 12 is a flowchart showing the steps of a process (that may be implemented as one or more software program or subprograms if the shown steps are divided) that, in part, determines the pre-determined parameter settings 105a and/or treatment time settings 105b (see steps 200, 220, 222, and 224).

The generator 104 (and/or the ablation probe unit 102) may be programmed by the operator and/or technicians at the laboratory and/or factory. The programming may be automatic or manual. "Programming" includes having the pre-determined settings 105 pre-entered and/or entering (inputting) the pre-determined settings 105 manually or automatically into the generator 104 (and/or the ablation probe unit 102) via operator input mechanisms. For example, the pre-determined settings 105 may be preprogrammed into an ablation probe unit 102, transmitted to the operator in the form of a programming signal (e.g. over the internet to be downloaded and installed in the ablation probe unit 102 or the generator 104), provided in the form of computer-readable media (e.g. a disc or a solid state USB drive), and/or provided as data (or a code) that may be manually entered into the ablation probe unit 102 (or the generator 104). Ideally, whichever method of entering/programming the ablation probe unit 102 (or the generator 104) is used, operator error is considered and eliminated as much as possible and appropriate checks are used. Preprogramming and some of the other means for programming the ablation probe unit 102 (or the generator 104) with the pre-determined settings would help to eliminate operator input errors. Another example of means for eliminating errors is that even if the ablation probe unit 102 (or the generator 104) is preprogrammed by the laboratory, the pre-determined settings might be displayed to the user for independent "verification" as the user could notice variations from normal pre-determined settings (e.g. the literature provided might provide a range and the operator would notice if the provided pre-determined settings 105 fell outside of the range). Yet another example is that the pre-determined settings might be provided as a code that, when input, would only function if it corresponded with a logical setting (e.g. if the person's age was also input into the ablation probe unit 102 and the code was not a logical setting based on the age, the ablation probe unit 102 would not function).

The pre-determined settings 105 for each TBA site may be included in the TBA surgical kit as a print out, on a disk or other computer readable storage media, or with instructions on how to obtain or download the information.

The pre-determined ablation means parameter settings 105a can also be referred to as "parameter settings 105a," "preferred parameter settings 105a," "optimal parameter settings 105a," "ideal parameter settings 105a," "pre-determined parameter settings 105a," "recommended parameter settings 105a," or "prescribed parameter settings 105a."

Tissue Trocar 146

If the ablation probe tip 108 is not self-introducing, at least one sharp instrument (that is preferably disposable) such as a tissue trocar 146 (and sometimes a plurality of tissue trocars) may be used by the operator to introduce (initially create) the access opening through the thick attached gingival tissue 122 that overlays third molar tooth buds 120. The tissue trocar tips are preferably sharp enough to be pushed and/or punched through the gingival tissue 122 into the base of the tooth bud. The diameter of the tissue trocar 146 rapidly increases up to 100% of the size of the ablation probe tip 108. After the tissue trocar 146 has created

the access opening, the tissue trocar 146 is removed and the ablation probe tip 108 is immediately placed into the access opening.

TBA Surgical Kit

The TBA surgical kit is a package that includes the majority of the necessary components and information for the surgical phase 90 of the TBA procedure 70. The TBA kit will be assembled (or the assembly will be completed) based on the patient's impressions and volume scans. Preferably, the TBA surgical kit has attractive packaging.

An exemplary TBA surgical kit may consist of (a) a custom surgical stent 110 for each arch as required, (b) at least one ablation probe tip 108 labeled its respective surgical site, (c) at least one tissue trocar 146 (if necessary), and (d) pre-determined settings 105 for each TBA site along with patient and operator identification.

If feedback controls are a part of the ablation probe tip design, then the correct in situ tissue temperature settings are preferably computed and supplied with the ablation probe tips 108 as part of the surgical kit.

The generator 104 and/or the hand pieces 106 are standard equipment in a dental office and/or can be purchased separately.

The ablation probe tips 108 may be pre-purchased (or extras may be kept in a practitioner's office) in which case the TBA surgical kit would provide a part number or other identifying information so that the practitioner would know which ablation probe tip 108 should be used with each guide 112.

It should be noted some of the components may not be part of the physical TBA surgical kit. For example, the pre-determined settings 105 may be provided electronically.

The TBA Procedure 70

Using the TBA procedure 70 described herein, the effective center of ablation 130a of the ablation probe tip 108 can be positioned at a pre-defined angle (ϕ) and pre-defined depth (x) so that the ablation probe tip's effective center of ablation 130a is positioned substantially in the "middle" of the tooth bud 130b within approximately 50%, 25%, or even less than 10% of the average diameter of the tooth bud 120. This is extremely accurate as compared to previous procedures.

FIG. 4 shows the steps and/or phases in an exemplary preferred TBA procedure 70: (1) routine screening and diagnosis 72; (2) pre-surgical scanning 82 (including taking impressions 84 and using scanning technology 86); (3) assembling a TBA surgical kit 88 (including pre-determined settings 105 and a stent 110); (4) operator delivery of the surgical phase 90 of the TBA procedure 70 (shown in more detail in FIG. 11); and (5) post-surgical (follow-up) steps 98. Steps (2) and (3) are also referred to jointly as the pre-surgical phase 80 during which steps are taken to create (including calculating, manufacturing, fabricating, selecting, and/or assembling) components of the TBA system 100 and/or the TBA surgical kit to be provided to the operator. Step (4) is also referred to as the surgical phase 90 of the TBA procedure 70 during which the steps shown in FIG. 11 are taken to ablate tooth buds 120.

(1) Screening Phase 72

Routine screening using panoramic or intra-oral X-ray imaging techniques is necessary to identify the presence of forming tooth buds 120 starting at age 6 through age 12 because of the wide range of ages involved with the formation of third molar tooth buds 120.

(2) Impressions and Scanning of Pre-Surgical Phase **80**

Once third molar tooth buds **120** have been identified to be present using standard screening methods (screening phase **72**), the next step is to pre-operatively measure the precise three-dimensional location and volume of each third molar tooth bud **120**. This may be practically accomplished using scanning technology **86** (e.g. computed tomography volume scanning such as CBCT). Scanning technology **86** can be used to accurately generate the necessary three-dimensional volume scans (computed tomography volume scans) and measurements ± 0.2 mm using, for example, the distal side of erupted first molars as durable physical landmarks (although it is possible to use soft tissue over bone as landmarks). The scanning technology **86** produces tooth bud size and position data **86'** (also referred to as "volume scans" and/or "measurements") that is provided for the step of producing the TBA surgical kit **88**. The tooth bud size and position data **86'** may be provided as a scanning technology file that can be any data file generated by the scanning technology **86** with the data necessary to manufacture or fabricate a stent **110**. One exemplary type of scanning technology file is a three-dimensional computer aided design (CAD) file.

An impression **84** of the patient's teeth and gum tissue (gingival tissue **122**) is made using standard impression materials such as PVS-type impression material (although other impression materials can be used). The impressions **84** are then processed and/or scanned using scanning technology (e.g. CBCT imaging by dentists and/or CT imaging in the laboratory), and the resulting volume scan of the impression is emailed (or otherwise transmitted or delivered) to a laboratory and/or factory where the volume scan is used for manufacturing or fabricating. It is still possible to physically mail the PVS dental impressions **84** to the designated laboratory and/or factory for manufacturing or fabricating.

Although the scanning technology is discussed primarily in terms of computed tomography volume scanning (e.g. CBCT technology), alternative scanning technologies including, but not limited to, ultrasound scanning technologies and future developed scanning technologies are included in the scope of the invention. Specialty software or programs may be used with the scanning technology **86** to accomplish the purpose described herein. It should be noted that alternative scanning technology **86** (including future developed scanning technology) may be used if it is able to accurately generate the necessary three-dimensional volume scans and measurements ± 0.2 mm using the distal side of erupted first molars (or other landmarks) as durable physical landmarks. It should be noted that alternative scanning technology (including future developed scanning technology) may also be used as long as two- or three-dimensional scanning results in the positioning of the effective center of ablation **130a** within approximately 50%, 25%, or even less than 10% of the average diameter of the tooth bud **120**.

(3) Assembling a TBA Surgical Kit **88**

The pre-surgical phase **80** of the TBA procedure **70** includes assembling a TBA surgical kit **88**. This step of assembling a TBA surgical kit **88** preferably includes computing pre-determined settings **105** and manufacturing or fabricating the stent **110** based on tooth bud size and position data **86'** obtained from the scanning technology **86**. The process of computing pre-determined settings **105** may be controlled by a process (that may be implemented by software or a program). The process of manufacturing or fabricating the stent **110** may also be controlled by a process (that may be implemented by software or a program).

After the impressions **84** are processed and/or scanned and the tooth bud size and position data **86'** is obtained, the process of manufacturing or fabricating the stent **110** may be carried out using direct-digital manufacturing or fabricating techniques similar to the processes used for manufacturing or fabricating implant surgical stents directly from CBCT scans (e.g. the processes used for fabricating SurgiGuide™ and other implant surgical guides) and the process used for manufacturing or fabricating orthodontic aligners (e.g. orthodontic aligners made by Align Technology or ClearCorrect). The direct-digital manufacturing or fabricating techniques, however, use the tooth bud size and position data **86'** to position and angle the surgical guides **112** on the distal aspects of the surgical stents **110** and use the erupted first molars as the primary landmark for positioning. Although manufacturing or fabricating will usually be done remotely in a laboratory and/or factory, it is possible that larger clinics will have the ability to manufacture or fabricate surgical stents **110** in their own in-house laboratory and/or factory.

Direct-digital manufacturing or fabricating techniques can be defined as any manufacturing or fabricating process that creates physical parts directly from data (e.g. three-dimensional CAD files) using manufacturing or fabricating techniques including, but not limited to, surgical stent manufacturing or fabricating technologies, rapid turn-around fabrication technologies, computer aided manufacturing (CAM), technologies using CAD, computer numerical control (CNC) milling, "additive" manufacturing, direct-digital laser stereolithography fabrication, "3-D printing," or any other manufacturing or fabricating means known or yet to be discovered that is capable of using the results generated by scanning to manufacture or fabricate the custom surgical stents. Because of the possibility for the integrated use of direct-digital volume scanning of impressions, low manufacturing costs, and rapid turn around times, use of direct-digital manufacturing or fabricating techniques is one preferred manufacturing or fabricating technique, but more traditional manufacturing or fabricating techniques that require more labor intensive manual laboratory processing could also be used.

At least one process that may be implemented as software or as at least one program (e.g. custom software enhancements in the CBCT software) will preferably assist in the direct-digital manufacturing or fabricating of the surgical stents **110** and define (and/or compute or calculate) the pre-determined settings **105**. This process would include defining (and/or computing or calculating) positioning and entry angle data required for placement of the ablation probe tip's effective center of ablation **130a** into the middle of the targeted tooth bud **120**. Additionally, tooth bud volumes are preferably computed (possibly using the scanning technology) and then the tooth bud volumes are used to determine the pre-determined settings **105** necessary to effect therapeutic ablation. Tooth bud volumes will generally range from 4.0 mm to 12.0 mm in diameter at ages 6-12. The ablation means **104'** and treatment times are preferably considered in the calculations. Companies that make CBCT imaging equipment promote the development of procedure-specific software in order to gain end-user acceptance of their imaging systems in the market place. The process may use calculations and/or look-up charts (e.g. based on experimental data) for determining the necessary settings.

FIG. **12** is a flowchart showing the steps of a process (that may be implemented as one or more software programs or subprograms if the shown steps are divided) for manufacturing or fabricating custom surgical stents **110** and/or determining the pre-determined parameter settings **105a**

and/or treatment time settings **105b**. As shown, the process begins with receiving pre-operative measurements of the precise three-dimensional location and volume of each third molar tooth bud and information regarding the ablation probe unit including its ablation means capabilities **200**. To make the stents **110**, the process would preferably include the following steps: (a) determining an entry point for an ablation probe tip **210**; computing the angle and depth of the path between the entry point and the middle of a tooth bud **212**; (b) taking into consideration the depth of the path, creating or selecting an ablation probe tip having the proper distance between its mechanical stop and its center of ablation so that the ablation probe tip will be inserted so that its center of ablation will be in the middle of the tooth bud **214**; (c) taking into consideration the angle and depth of the path and the thickness of the surgical stent, computing the surgical guide pathway through which the ablation probe tip will be inserted so that its center of ablation will be in the middle of the tooth bud **216**; and (d) providing the surgical guide pathway as output for the creation of a surgical stent with surgical guides **218**. To calculate the pre-determined parameter settings **105a** and/or treatment time settings **105b**, the process would preferably include the following steps: (a) taking into consideration the information regarding the ablation probe unit including its ablation means capabilities, determining the proper power settings **220**; (b) taking into consideration the information regarding the ablation probe unit including its ablation means capabilities, determining the proper time settings **222**; and (c) providing the proper power and time settings as output for use in programming the ablation probe unit or generator **224**.

As described above, in addition to the surgical stent(s) **110** and the pre-determined settings **105**, the TBA surgical kit may include at least one ablation probe tip **108** labeled for its respective surgical site, at least one tissue trocar **146** (if necessary), and patient and operator identification.

The TBA surgical kit is provided to the operator.

(4) Surgical Phase **90**

FIGS. **6-10** show graphically, and FIG. **11** shows as a flow chart, the surgical phase **90** of the TBA procedure. The surgical phase **90** may be performed by a dental operator (dental practitioner) in his office (e.g. a pediatric office and/or general dental office) under normal office conditions. At this point, the generator **104** has been programmed with the pre-determined settings **105** and normal surgical procedures have been followed. The generator **104** is preferably tuned so that the ablation means **104'** is set to ablate the small, substantially spherical ablation volumes of third molar tooth buds **120** in order to minimize (or possibly eliminate) collateral osseous and soft tissue damage, especially damage to adjacent second molars that are likely not yet erupted. Further, the surgical phase **90** uses single-use and disposable delivery systems that use components designed for intra-oral use.

Summarily, as shown in FIG. **11**, the first step is physically seating a surgical stent **160** in a patient's mouth. Next, the operator makes an access path at the at least one tooth bud surgical site **162**. The operator also places the ablation probe tip so that the center of ablation is in the middle of a tooth bud at the at least one tooth bud surgical site (using the custom surgical stent to guide the placement) **164**. It should be noted that if the ablation probe tip is "self-introducing," the step of making an access path and the step of placing the ablation probe tip may occur simultaneously. Then, the at least one tooth bud is at least partially ablated **166** and the

ablation probe tip is removed from the tooth bud **168**. These and other exemplary steps are detailed in the following paragraphs.

The operator preferably starts the surgical phase **90** by placing the surgical stent **110** into place onto the patient's teeth prior to administering local anesthetic to the surgical site. The local anesthetic will then be administered through the surgical stent **110** and guides **112** that are in close approximation with the gingival tissue **122**, thus reducing the amount of anesthetic necessary because of the precise placement of anesthetic agent. Achieving local anesthesia in this procedure will be easier than anesthetizing lower permanent molar teeth for routine fillings since only soft tissues, which will be 8.0 mm to 15.0 mm deep, are involved.

The step of physically seating a surgical stent **110** may also include physically seating the surgical stent in a patient's mouth, physically seating the surgical stent on a patient's erupted teeth, physically seating the surgical stent on at least one tooth in a patient's mouth, physically seating the surgical stent on a patient's soft tissue, physically seating the surgical stent on a patient's bone, or a combination of the above steps (e.g. physically seating the surgical stent on a patient's teeth, soft tissue, and bone).

Once the custom surgical stent **110** is in place and the patient is fully anesthetized, the operator then mechanically gains access to the tooth bud **120** through the stent surgical guides **112** by creating (introducing) a small surgical access path opening through the gingival tissue **122** approximately 0.1 mm to 2.0 mm (and more particularly 0.5 mm to 1.0 mm) in diameter using tissue trocars. If the ablation probe tips **108** are designed to be strong enough and sharp enough to act as "self-introducing" probe tips, they can be used to introduce the surgical access path. On the other hand, if the ablation probe tip itself is not self-introducing, the surgical access path may be introduced using a then there will be no need for separate tissue trocar **146**.

It should be noted that the surgical access path is preferably an incision, a puncture, or a hole through the gingival tissue **122**. If a self-introducing probe tip is used, the surgical access path has substantially the same diameter as the ablation probe tip **108**. If the probe tip is not self-introducing, the surgical access path may be a sutureless puncture (0.1 mm to 2.0 mm in diameter) or, more particularly, a sutureless puncture (0.5 mm to 1.0 mm in diameter). Alternatively, a trocar "punch" may be made through tough gingival tissue **122**. Regardless of the procedure used to introduce the surgical access path, using a surgical access path to gain access or allow placement of the ablation probe tips **108** to the tooth bud **120** does not kill, damage, or otherwise cause necrosis to the surrounding soft tissues (e.g. gingival tissues **122**). This can be compared to other processes such as coring, boring, cutting, electrourge ablating, or other invasive procedures that kill, damage, and/or otherwise cause necrosis to the soft tissue to which the invasive procedure has been applied. Although the preferred procedures for introducing the surgical access path might kill individual cells, the soft tissue (the gingival tissue **122**) does not become necrosed because the tissue is a collection of cells that can heal itself.

As shown in FIGS. **6** and **7**, the next step in the surgical phase **90** is to insert the designated ablation probe tip **108** through the surgical stent **110** and into the tooth bud space until it is mechanically "stopped" in order to position the probe to the prescribed depth (which would be the pre-defined depth). The surgical stent **110** and its surgical guides **112** are used to control the angle (ϕ) and depth (x) of the

ablation probe tip **108** so that the effective center of ablation **130a** of the ablation probe tip is in the middle of the tooth bud **130b**. It should be noted that the effective center of ablation **130a** for any given ablation technology does not necessarily correspond with the tip of the ablation probe. For instance, microwave ablation probes have windows or slots that may be 0.5 mm to 2.0 mm from the tip depending on the frequency of the wavelength used. Cryoablation probes have their center of ablation roughly in the middle of the probe, depending on the design and refrigerant used. A mechanical stop structure **140** on the ablation probe tip **108** preferably seats firmly onto the mechanical stop structure **142** of the surgical stent guide **112** to prevent over extension of the ablation probe tip **108**.

FIG. **8** shows embedded temperature sensors **144** (or other types of feedback control mechanisms) that may be used during the ablation process. An independent feedback process using the temperature sensors **144** is preferable for this clinical procedure. Use of temperature sensors **144** along with monitoring probe impedance characteristics and percentage of reflected energy in RF/MW circuits will provide “go/no go” output for the clinician. Control algorithms are preferably used to accelerate initial ablation means **104'** input followed by lower-level temperature maintenance for a defined period of time with independent confirmation that results in a fast process while simultaneously assuring complete tooth bud ablation.

FIG. **9** shows the actual ablation process. Activation of the ablation probe unit **102** to perform the ablation process is executed according to the pre-determined settings **105**. Activation of the ablation probe unit **102** causes the generator **104** to provide the ablation means **104'** that passes through the hand piece **106** and the ablation probe tip **108** and into the tooth bud **120**. This step of at least partially ablating the tooth bud is preferably accomplished without ablating any surrounding gingival tissue (although a minimal amount of surrounding gingival tissue may be ablated as an accidental byproduct of the step). This can also be thought of as the activation of the ablation probe unit **102** creating a zone of ablation that resides predominantly or completely within the tooth bud **120**. The feedback control mechanisms **144** assure successful delivery of adequate ablation means **104'** to ablate the tooth bud **120** while minimizing damage to adjacent osseous and soft tissues by, for example, eliminating overheating. Given the small tissue volumes involved for pediatric patients, activation using an RF ablation means **104'** would have an ablation time that is preferably less than three (3) minutes and activation using an MW ablation means **104'** would have an ablation time that is preferably less than thirty (30) seconds.

FIG. **10** shows the ablation probe tip **108** being removed from the now ablated tooth bud **120'**. As shown in this figure, any access path created by the procedure rapidly closes.

(5) Post-Surgical Phase **98**:

After the surgical phase **90**, the patient may have follow-up including, but not limited to, post-surgical instructions and, if necessary follow-up care and screening.

Post-surgical instructions that may be given to parents includes the following: kids can go out and play immediately unless they were sedated, no post-surgical pain medication is necessary, bleeding (if any) will be gone in minutes, and post-surgical X-ray screening may be necessary at patient's next routine 6-month hygiene cleaning appointment to verify full ablation.

Simulated TBA Procedure **70**

The following paragraphs, along with FIGS. **13-29**, detail an exemplary simulated TBA procedure **70** including routine

screening and diagnosis **72**, the pre-surgical phase **80**, and the surgical phase **90**. In several of these figures, a patient's mouth **124** (with gums **122** and teeth **126**) is shown that looks like a stone model, but it should be understood that unless otherwise specified the shown mouth **124** would be a live patient's mouth.

As shown in FIG. **4**, the TBA procedure begins with routine screening and diagnosis **72**. FIG. **13** is a panoramic X-ray showing a patient whose third molar tooth buds **120** in the #17 & #32 positions are treatable by a TBA procedure **70**. FIG. **14** is a pre-operative CBCT scan (although other types of volume scanning could be used) of a patient. In a real procedure, the volume scan would be taken of the specific patient on whom the TBA procedure **70** is being performed. This CBCT “reconstructed” panoramic scan has a 1.0 mm scale along its bottom edge. FIG. **15** is a series of CBCT volume scan cross-sections showing successive 1.0 mm slices through both #17 and #32 in 1.0 mm increments. Each X-ray corresponds to 1.0 mm locations along the scale of FIG. **14**. The left-side scale is 1.0 mm vertically. The maximum tooth bud diameters are measured to be 8.0-9.0 mm.

FIG. **16** shows a pre-operative upper-arch impression **84** being taken of the simulated patient's mouth **124** (shown as a stone model for clarity, but an impression **84** would be taken of the patient himself) using an impression tray **128**. It is assumed that all four tooth buds of the wisdom teeth are present in the simulated patient. FIG. **17** is a cross-sectional view of the upper-arch impression **84** being taken of a simulated patient. FIG. **18** shows the completed upper-arch impression **84**. A similar process would be performed to manufacture or fabricate a pre-operative lower-arch impression **84**. At this time the practitioner may send impressions **84** and volume scan data to a laboratory and/or factory for processing.

The laboratory and/or factory uses the impressions **84** and volume scan data (scanning technology file) to create (including calculating, manufacturing, fabricating, selecting, and/or assembling) components of the TBA system **100** (including the surgical stents **110** and the pre-determined settings **105**). The surgical stents **110** and the pre-determined settings **105** and other components are then assembled into the TBA surgical kit to be provided to the operator.

FIG. **19** shows the completed upper-arch impression **84**, along with a stone model **85** that will serve as a “positive” for manufacturing or fabricating a surgical stent **110** for that patient's upper-arch. Alternatively, when using stereolithography manufacturing to manufacture or fabricate surgical stents **110**, the impressions **84** can be computed tomography (“CT”) scanned to digitize as an alternative to making physical intermediates. The CT volume scan file (scanning technology file) can then be emailed (or otherwise directly transmitted) for direct manufacturing or fabricating. Alternatively, the practitioner may handle the processing in-house.

FIG. **20** is a CBCT scan with notations showing the measurement of the perpendicular angle of entry into the tooth bud **120**. The measurement is based on the distal aspect of the molar and the occlusal bite plane of the teeth. FIG. **21** is a series of X-rays with notations showing the measurement of the lateral angle of entry. The measurement is determined relative to the vertical axis in order to avoid the jaw's boney interferences during surgical placement of the ablation probe unit **102**. FIG. **22** is a CBCT scan with highlights showing the computed volume of each tooth bud **120**. CBCT volume data is used to determine and/or calculate the pre-determined settings **105**.

FIG. 23 shows the resulting surgical stent 110 that will be placed in a patient's mouth 124. The shown stent has two surgical guides 112 based upon the location of the patient's two tooth buds to be ablated.

The surgical stent(s) 110 and the pre-determined setting(s) 105 are provided to the operator along with the rest of the TBA surgical kit.

Prior to the surgical phase 90 of the TBA procedure 70, the ablation probe unit 102 and/or the generator 104 should be set up so that at least one pre-determined setting 105 is correctly entered for at least one tooth bud 120 with safety interlocks carefully considered. (The pre-determined settings 105 may all be entered prior to the surgical phase 90 or they may be entered one at a time.) The surgical phase 90 of the TBA procedure 70 may then be performed.

FIG. 24 shows topical anesthetic 87 being applied to the base of the surgical guide 112 (FIG. 24) prior to the surgical stents 110 being seated in a patient's mouth 124.

FIG. 25 shows the surgical stent 110 being seated on the upper-arch of the simulated patient's mouth 124 (shown as a stone model for clarity). This process would be repeated on the lower-arch of the simulated patient.

FIG. 26 shows a local anesthetic being injected 89 into each site through a surgical guide 112 of the stent 110.

FIG. 27 shows a tissue trocar 146 being used to create an access path through the gingival tissue 122 to the base of each tooth bud 120. The tissue trocar 146 is only necessary if self-introducing ablation probe tips 108 are not used.

FIG. 28 shows an ablation probe tip 108 with mechanical stop structure 140' (shown as a shoulder) being inserted through the surgical guide 112. This would be similar to the position of the ablation probe tip 108 in FIG. 6.

FIG. 29 shows the ablation probe tip 108 positioned through the surgical guide 112 and into the tooth bud 120 through the surgical guide 112 so that the ablation probe tip's effective center of ablation 130a is in the middle of each tooth bud 120. This would be similar to the position of the ablation probe tip 108 in FIG. 7.

The ablation means 104' is delivered in this position (FIG. 9). The ablation means 104' is delivered based on the pre-determined settings 105 (e.g. times, intensities, and other prescribed settings unique to each tooth bud).

The ablation probe tip 108 would then be removed and the process repeated at the site of each tooth bud 120. Once the entire surgical phase 90 is complete, the surgical stents 110 are removed.

Finally, the dental practitioner or an assistant provides post-surgical instructions to the patient or a caregiver of the patient.

Alternative Scanning and Fabrication of Custom TBA Surgical Kits

An alternative to the pre-surgical phase 80 of the TBA procedure 70 described above includes simultaneous three-dimensional scanning of both hard tissues (bone and teeth) and soft tissues (tooth bud 120 and gingival tissue 122). From the information obtained using this unique simultaneous three-dimensional scanning, a custom surgical stent 110 may be manufactured or fabricated. As discussed, the custom surgical stent 110 is used in the surgical phase 90 to help with the placement of the center of ablation 130a into a tooth bud 120 that results in tooth agenesis.

The simultaneous three-dimensional scanning uses a single scan to obtain both soft tissue and hard tissue information. Soft tissue information generally does not show on a scan, although progress in volume scanning is improving and this may be possible in the near future. Known and future technologies able to provide a scan image of soft

tissue are included in the scope of this invention. A typical X-ray scan will only show the hard tissue. So to obtain both soft and hard tissue information using simultaneous three-dimensional scanning, a dental impression 84 is used that can be viewed on an X-ray. The dental impression 84 is made of materials that are preferably "contrast optimized" for high resolution X-ray volume scanning. The ideal level of contrast agent in the range of 25% to 75% radiopacity (such as barium or iodine based compounds) is mixed into the dental impression materials so that the highest level of surface detail can be picked up on when volume scanning the dental impression 84. The dental impression 84 is placed in the patient's mouth 124 during the X-ray volume scan. The resulting X-ray volume scan image would show the tooth distinguished (is visible) and the dental impression 84 distinguished (is visible) and the void therebetween would be the soft tissue and would therefore be "visible." The resulting X-ray volume scan with both hard and soft tissue information may then be used to formulate the custom stent 110 used in the surgical phase 90 described herein. In other words, an X-ray volume scan image is generated in which hard tissue (e.g. a tooth) is visible hard tissue and the dental impression 84 is a visible dental impression and soft tissue (e.g. gingival tissue 122) is "visible" as the space between the visible hard tissue and the visible dental impression.

One separate preferred pre-surgical phase 80 of the TBA procedure 70 preferably includes using X-ray volume scans of dental impressions 84 to manufacture or fabricate surgical stents 110. The X-ray volume scan of the dental impression 84 is "super imposed" over the patient X-ray volume scan (e.g. CBCT scanning) using the dental hard tissues (the teeth) to "snap" the two volume scans together into an accurate overlay so that soft tissues of the mouth (which cannot be X-ray volume scanned directly) are accurately defined for the surgical stent manufacturing or fabricating (which must take into account the soft tissue and teeth) and probe positioning (which must take into account the tooth bud positioning from the patient's CBCT scan).

One separate preferred pre-surgical phase 80 of the TBA procedure 70 preferably includes using dental impression materials that are "contrast optimized" for high resolution X-ray volume scanning that is then used to manufacture or fabricate surgical stents 110. The ideal level of contrast agent (such as barium or iodine based compounds) is mixed into the dental impression materials so that the highest level of surface detail can be picked up on when CT volume scanning the dental impression 84.

Alternative Procedures and Systems

Separate preferred surgical procedures preferably include the ablation of "non-tooth" bud lesions or tumors of the maxilla or mandible. In such a situation, a custom stent would be manufactured or fabricated with guides to guide an ablation probe tip 108 to such a lesion or tumor located at least one lesion or tumor surgical site. The process could then be used to ablate such lesion or tumor.

Separate TBA surgical procedures preferably include the use of ultrasound scanning with combined ultra-high energy ultrasound ablation but without the use of a surgical stent for transgingival tooth bud ablation that results in tooth agenesis. This can be described as direct ultrasound scanning with ultra-high energy ultrasound built into the same scanning head.

Comparison to the Silvestri Study

As set forth in the Background section of this document, the article entitled "Selectively Preventing Development Of Third Molars In Rats Using Electrosurgical Energy" by Silvestri et al. describes a pilot study that tests the hypothesis

that third molars can be selectively prevented from developing. The results of the Silvestri study were mixed at best, with only ten rats out of thirty-three showing the desired result of no intraoral or radiographic evidence of third molar development. One reason that the Silvestri process was not successful may have had to do with the fact that the Silvestri process was inexact. For example, the Silvestri process relies on molds taken from molds of the mouths of euthanized rat pups rather than using molds fabricated for the rat pup on which the procedure was to be performed. The present invention uses the patient's mouth on which the procedure is to be performed. Another way in which the Silvestri process was inexact was that the Silvestri process did not locate the forming tooth bud **120**. More specifically, the Silvestri process did not locate or determine the location of the forming tooth bud **120** pre-operatively relative to the landmarks that he used. Silvestri even states ". . . when electrosurgical energy is applied near the invisible tooth anlage in the tiny mouth of newborn rats, the effects of the electrosurgical energy cannot be nearly as local or precise. The embryonic tooth-forming tissues of the third molar [lay] fractions of a millimeter below the oral mucosa and cannot be seen. As a result, it was not possible to predictably protect and isolate the vulnerable developing bone from the energy and heat of the electrosurgical energy. The result was a relatively large, unpredictable area of tissue damage during treatment and a wide range of bony developmental effects seen after the rats were euthanized." The TBA procedure **70** described herein can be distinguished from the Silvestri procedure in several ways including for example, that (1) the TBA procedure **70** described herein is a minimally invasive procedure consisting of introducing a surgical access path at each tooth bud surgical site as opposed to the boring, killing, and damaging procedure described by Silvestri, (2) the TBA procedure **70** described herein is performed in such a manner that it can be described as exact (e.g. using the patient's mouth as the mold for manufacturing or fabricating the surgical stent **110**, taking exact measurements of the patient's mouth (including the position of the tooth bud **120**), and using calculated parameter and time settings **105b**) as opposed to the Silvestri procedure that can be described as inexact, and (3) the TBA procedure **70** described herein can predictably ablate tooth buds **120** as opposed to the Silvestri procedure that was essentially unpredictable and could never, under any circumstances, be considered for treating human patients.

Flow Charts

FIGS. **4**, **11**, and **12** are flow charts illustrating processes, methods, and/or systems. It will be understood that at least some of the blocks of these flow charts, components of all or some of the blocks of these flow charts, and/or combinations of blocks in these flow charts, may be implemented by software (e.g. coding, software, computer program instructions, software programs, subprograms, or other series of computer-executable or processor-executable instructions), by hardware (e.g. processors, memory), by firmware, and/or a combination of these forms. As an example, in the case of software, computer program instructions (computer-readable program code) may be loaded onto a computer (or on a special purpose machine such as a volume scanner or scanning technology) to produce a machine, such that the instructions that execute on the computer create structures for implementing the functions specified in the flow chart block or blocks. These computer program instructions may also be stored in a memory that can direct a computer to function in a particular manner, such that the instructions stored in the memory produce an

article of manufacture including instruction structures that implement the function specified in the flow chart block or blocks. The computer program instructions may also be loaded onto a computer (or on a special purpose machine such as a volume scanner or scanning technology) to cause a series of operational steps to be performed on or by the computer to produce a computer implemented process such that the instructions that execute on the computer provide steps for implementing the functions specified in the flow chart block or blocks. The term "loaded onto a computer" also includes being loaded into the memory of the computer or a memory associated with or accessible by the computer (or on a special purpose machine such as a volume scanner or scanning technology). The term "memory" is defined to include any type of computer (or other technology)-readable media including, but not limited to, attached storage media (e.g. hard disk drives, network disk drives, servers), internal storage media (e.g. RAM, ROM), removable storage media (e.g. CDs, DVDs, flash drives, memory cards, floppy disks), and/or other storage media known or yet to be discovered. The term "computer" is meant to include any type of processor, programmable logic device, or other type of programmable apparatus known or yet to be discovered. Accordingly, blocks of the flow charts support combinations of steps, structures, and/or modules for performing the specified functions. It will also be understood that each block of the flow charts, and combinations of blocks in the flow charts, may be divided and/or joined with other blocks of the flow charts without affecting the scope of the invention. This may result, for example, in computer-readable program code being stored in whole on a single memory, or various components of computer-readable program code being stored on more than one memory.

ADDITIONAL INFORMATION

It is to be understood that the inventions, examples, and embodiments described herein are not limited to particularly exemplified materials, methods, and/or structures. Further, all publications, patents, and patent applications cited herein, whether supra or infra, are hereby incorporated by reference in their entirety.

Please note that the terms and phrases may have additional definitions and/or examples throughout the specification. Where otherwise not specifically defined, words, phrases, and acronyms are given their ordinary meaning in the art. The following paragraphs provide some of the definitions for terms and phrases used herein.

The terms "fabricating" and/or "manufacturing" include any suitable means of making a component (e.g. stent **110**). Although the terms are used together throughout most of the specification (e.g. "manufacturing or fabricating"), the absence of one term or another is irrelevant because they are used herein synonymously.

The terms "proper," "correct," "optimal," and "ideal," are relative and may become more accurate as technology is developed. For example, when used in terms of the pre-defined angle (ϕ) and pre-defined depth (x) that are calculated and/or prescribed (e.g. the "proper angle and depth," the "correct angle and depth," the "optimal angle and depth," or the "ideal angle and depth"), these phrases are meant to include the best possible angle and depth that is calculated using the best available information and technology.

The terms "provide" and "providing" (and variations thereof) are meant to include standard means of provision including "transmit" and "transmitting," but can

also be used for non-traditional provisions as long as the data is "received" (which can also mean obtained). The terms "transmit" and "transmitting" (and variations thereof) are meant to include standard means of transmission, but can also be used for non-traditional transmissions as long as the data is "sent." The terms "receive" and "receiving" (and variations thereof) are meant to include standard means of reception, but can also be used for non-traditional methods of obtaining as long as the data is "obtained."

It should be noted that the terms "may" and "might" are used to indicate alternatives and optional features and only should be construed as a limitation if specifically included in the claims. It should be noted that the various components, features, steps, phases, or embodiments thereof are all "preferred" whether or not it is specifically indicated. Claims not including a specific limitation should not be construed to include that limitation.

It should be noted that, unless otherwise specified, the term "or" is used in its nonexclusive form (e.g. "A or B" includes A, B, A and B, or any combination thereof, but it would not have to include all of these possibilities). It should be noted that, unless otherwise specified, "and/or" is used similarly (e.g. "A and/or B" includes A, B, A and B, or any combination thereof, but it would not have to include all of these possibilities). It should be noted that, unless otherwise specified, the term "includes" means "comprises" (e.g. a device that includes or comprises A and B contains A and B but optionally may contain C or additional components other than A and B). It should be noted that, unless otherwise specified, the singular forms "a," "an," and "the" refer to one or more than one, unless the context clearly dictates otherwise.

The terms and expressions that have been employed in the foregoing specification are used as terms of description and not of limitation, and are not intended to exclude equivalents of the features shown and described. This application is intended to cover any adaptations or variations of the present invention. It will be appreciated by those of ordinary skill in the art that any arrangement that is calculated to achieve the same purpose may be substituted for the specific embodiment shown. It is also to be understood that the following claims are intended to cover all of the generic and specific features of the invention herein described and all statements of the scope of the invention which, as a matter of language, might be said to fall therebetween.

What is claimed is:

1. A custom surgical stent configured for use in a tooth bud ablation procedure that results in tooth agenesis, the custom surgical stent for use with an ablation probe tip having a center of ablation and mechanical stop structure, said stent comprising:

- (a) a custom surgical stent with at least one surgical guide corresponding to at least one tooth bud surgical site;
- (b) said at least one surgical guide having guiding structure to guide placement of an ablation probe tip at a

pre-defined angle so that a center of ablation of said ablation probe tip is in the middle of a tooth bud;

- (c) said at least one surgical guide having mechanical stop structure to limit the depth of said ablation probe tip to a pre-defined depth; and
- (d) said mechanical stop structure of said at least one surgical guide and said mechanical stop structure of said ablation probe tip together limiting the depth of said ablation probe tip through said at least one surgical guide until there is a mechanical stop between said mechanical stop structure of said at least one surgical guide and said mechanical stop structure of said ablation probe tip;
- (e) wherein said mechanical stop structure of said at least one surgical guide and said guiding structure of said at least one surgical guide together position said ablation probe tip so that said center of ablation is positioned within less than 50% of the average diameter of the tooth bud.

2. The stent of claim 1, wherein said at least one surgical guide is directly associated with said custom surgical stent.

3. The stent of claim 1, wherein the combination of said custom surgical stent and said at least one surgical guide is one piece.

4. A custom surgical stent configured for use in a tooth bud ablation procedure that results in tooth agenesis, the custom surgical stent for use with an ablation probe tip having a center of ablation and mechanical stop structure, said stent comprising:

- (a) a custom surgical stent with at least one surgical guide corresponding to at least one tooth bud surgical site;
- (b) said at least one surgical guide having guiding structure to guide placement of an ablation probe tip at a pre-defined angle so that a center of ablation of said ablation probe tip is in the middle of a tooth bud;
- (c) said at least one surgical guide having mechanical stop structure to limit the depth of said ablation probe tip to a pre-defined depth;
- (d) said guiding structure to guide placement of the ablation probe tip to a computed pre-defined angle of ablation based on information obtained from a volume scan image; and
- (e) said mechanical stop structure of said at least one surgical guide limiting the depth of said ablation probe tip to a computed pre-defined depth of ablation based on information obtained from a volume scan image;
- (f) wherein said mechanical stop structure of said at least one surgical guide and said guiding structure of said at least one surgical guide position said ablation probe tip so that said center of ablation is positioned within less than 50% of the average diameter of the tooth bud.

5. The stent of claim 4, wherein said at least one surgical guide is directly associated with said custom surgical stent.

6. The stent of claim 4, wherein the combination of said custom surgical stent and said at least one surgical guide is one piece.

* * * * *